



Representation Stability and Finite Orthogonal Groups

Arun S. Kannan¹ · Zifan Wang¹

Received: 5 April 2022 / Accepted: 8 February 2023
© The Author(s) 2023

Abstract

In this paper, we prove homological stability results about orthogonal groups over finite commutative rings where 2 is a unit. Inspired by Putman and Sam (2017), we construct a category $\mathbf{OrI}(R)$ and prove a local Noetherianity theorem for the category of $\mathbf{OrI}(R)$ -modules. This implies an asymptotic structure theorem for orthogonal groups. In addition, we show general homological stability theorems for orthogonal groups, with both untwisted and twisted coefficients.

Keywords Representation stability · Homological stability · Finite orthogonal groups · Central stability

Mathematics Subject Classification (2010) 18A25 (primary) · 20C33 (secondary)

1 Introduction

1.1 A Motivating Example

Consider a compact oriented manifold X with nonempty boundary. The configuration space of n points in X is defined as the space of all possible choices of n distinct points in X :

$$\mathrm{Conf}_n(X) = X^n \setminus \{(x_1, x_2, \dots, x_n) \mid x_i = x_j \text{ for some } i \neq j\}.$$

There is a natural action of the symmetric group S_n on $\mathrm{Conf}_n(X)$ by permuting the coordinates. Note that there exist natural maps $\mathrm{Conf}_{n+1}(X) \rightarrow \mathrm{Conf}_n(X)$ by forgetting the

Presented by: Henning Krause

✉ Arun S. Kannan
akannan@mit.edu

Zifan Wang
atticusw@mit.com

¹ Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA

$(n + 1)$ th coordinate. These maps induce morphisms at each level of cohomology, with coefficients in a fixed field \mathbf{k} (or more generally a Noetherian ring) with $\text{char } \mathbf{k} = 0$:

$$H^i(\text{Conf}_n(X); \mathbf{k}) \rightarrow H^i(\text{Conf}_{n+1}(X); \mathbf{k}).$$

Denote by $\text{UConf}_n(X)$ the unordered configuration space, which is the orbit space of the action of S_n on $\text{Conf}_n(X)$. In the unordered case, it is a classical result [11] that for any fixed index i , $H^i(\text{UConf}_n(X); \mathbf{k}) \cong H^i(\text{UConf}_{n+1}(X); \mathbf{k})$ for all $n \gg 0$. For $\text{Conf}_n(X)$, however, this does not hold. Instead, we have *representation stability*: notice that the action of S_n on $\text{Conf}_n(X)$ induces a representation of S_n on $V_n := H^i(\text{Conf}_n(X); \mathbf{k})$, and therefore V_n splits into the direct sum of irreducible representations, which are naturally parametrized by partitions of n . For any partition λ of a positive integer k , we write $c_\lambda(V_n)$ to denote the multiplicity inside the expansion of V_n of the irreducible representation corresponding to the partition $(n - k, \lambda)$ of n . In [3], it was shown that there exists N such that the following three properties hold for all $n > N$:

- (1) Injectivity: the maps $V_n \rightarrow V_{n+1}$ are injective;
- (2) Surjectivity: the image of V_n spans V_{n+1} as a $\mathbf{k}[S_{n+1}]$ -module;
- (3) Multiplicity stability: for any k and any partition λ of k , $c_\lambda(V_n) = c_\lambda(V_{n+1})$.

A direct corollary is that the dimensions of V_n exhibit polynomial growth as $n \gg 0$.

1.2 Background, History, and Known Results

In general, the notion of representation stability considers a sequence of representations V_n of a family of groups G_n , with maps $V_n \rightarrow V_{n+1}$ and $G_n \rightarrow G_{n+1}$ between them that are compatible with the action of G_n on V_n . This framework was first introduced in [5] to describe the frequent observation that various representation-theoretic properties of V_n stabilize for sufficiently large n . Stability came to possess a very broad meaning: one example is homological stability, which holds for $\text{UConf}_n(X)$ as discussed above. Another example is multiplicity stability, which holds for $\text{Conf}_n(X)$. In Section 4 of this paper, we prove for orthogonal groups a categorical version of representation stability, phrased in terms of Kan extensions.

A framework was developed in [4], involving the functor category of modules over a category named **FI**. This category of **FI**-modules carries the rich structure of an abelian category, and it acts as a large algebraic structure which altogether controls the growth behavior of a sequence of representations, thus providing a fundamental explanation of various stability behavior associated with representations of the symmetric groups S_n .

The objects of the category **FI** are finite sets, and the morphisms between them are injections (hence the name **FI**). The reason that **FI** plays a prominent role in the representation stability of symmetric groups is because the automorphism group of an n -element object in **FI** is precisely S_n . Key to the utility of **FI**-modules is a certain Noetherian property that roughly states that any submodule of a finitely generated **FI**-module is also finitely generated. This local Noetherianity property is a key ingredient in [4]'s proof of representation stability of $\text{Conf}_n(X)$.

In [13], analogues (**VIC**-modules, **SI**-modules) to the functor category of **FI**-modules were constructed by replacing the symmetric groups with the general linear groups and the symplectic groups (over finite rings). Similar Noetherian properties and asymptotic structure theorems were proven, as well as broad homological stability theorems with twisted coefficients. Some of these results were strengthened in [12], where an explicit bound for the stability degree was shown.

1.3 Main Results

Our motivation for this paper was that among the classical groups, only the orthogonal groups were not studied much in the context of representation stability. The main obstacle in this case stems from the fact that, unlike alternating forms and symplectic groups, more than one isometry class of symmetric bilinear forms exists, even for modules over finite rings.

In this paper, we address this problem for orthogonal groups over finite commutative rings R where 2 is a unit. Extending methods in [13], we construct a category $\mathbf{OrI}(R)$ whose objects are finite-rank free R -modules equipped with a symmetric bilinear form, and morphisms are isometries. Associated to this category is the functor category $\mathbf{Rep}_{\mathbf{k}}(\mathbf{OrI}(R))$ whose objects are functors from $\mathbf{OrI}(R)$ to the category of \mathbf{k} -modules and morphisms are natural transformations (here \mathbf{k} is a Noetherian ring). Objects of this category are called $\mathbf{OrI}(R)$ -modules.

Our first result is that $\mathbf{Rep}_{\mathbf{k}}(\mathbf{OrI}(R))$ is locally Noetherian (Theorem 3.2), similar to the functor categories of \mathbf{FI} -modules, \mathbf{VIC} -modules, and \mathbf{SI} -modules, as discussed in the previous subsection. We then prove that this implies an asymptotic structure theorem in the sense of [13] for orthogonal groups (Theorems 4.5, 4.13 and Corollary 4.9). More explicitly, this means that for any finitely generated $\mathbf{OrI}(R)$ -module M and for any morphism $f : V \rightarrow W$ for $V, W \in \mathbf{OrI}(R)$ such that V has sufficiently large rank, we have:

- (1) Injective stability: $M(f)$ is injective;
- (2) Surjective stability: under the action of $M(\text{Aut}_A(W))$ the image of $M(f)$ generates $M(W)$;
- (3) Central stability: For sufficiently large N , M is the left Kan-extension of the restriction of M to the subcategory of objects whose rank is at most N .

Finally, we apply Theorem 3.2 to prove a general homological stability theorem with twisted coefficients for (indefinite) orthogonal groups over fields (Theorem 5.1). In particular, suppose R is a finite field with odd characteristic and V, W are vector spaces over R equipped with symmetric nondegenerate bilinear forms B, B' , respectively. Then, if $V \rightarrow W$ is an isometry, for each k there is an induced map

$$H_k(O(V, B); M(V, B)) \rightarrow H_k(O(W, B'); M(W, B'))$$

on the homologies of the corresponding orthogonal groups with coefficients over a finitely generated $\mathbf{OrI}(R)$ -module M over a prime field \mathbf{k} whose characteristic is coprime to that of R . This in particular implies that the homology with coefficients taken over \mathbf{k} stabilizes (Corollary 5.5).

Future directions to consider include applying the results in this paper to the context of congruence subgroups of orthogonal groups over number rings and also to mapping class groups of high-dimensional manifolds, as in [13].

1.4 Roadmap

This paper is structured as follows. In Section 2, we review the framework of functor categories of modules over a category, recall the definition of column-adapted maps, and recount the theory of symmetric bilinear forms over finite commutative rings. In Section 3, we define the category $\mathbf{OrI}(R)$ and the category of $\mathbf{OrI}(R)$ -modules, and show that the latter is locally Noetherian. In Section 4, we show the asymptotic structure theorem, which is a strong stability result for finitely generated $\mathbf{OrI}(R)$ -modules. Finally, in Section 5, we show

homological stability with twisted coefficients (coefficients determined by some finitely generated $\mathbf{OrI}(R)$ -module), as well as stability with untwisted coefficients.

2 Preliminaries

2.1 Finitely Generated \mathcal{C} -Modules

We begin by reviewing the necessary framework of functor categories of modules over a category.

Definition 2.1 Let \mathcal{C} be a category and \mathbf{k} a ring. A \mathcal{C} -module over \mathbf{k} is a functor $M : \mathcal{C} \rightarrow \mathbf{Mod}_{\mathbf{k}}$, where $\mathbf{Mod}_{\mathbf{k}}$ is the category of \mathbf{k} -modules. If the ring \mathbf{k} is clear from the context, we shall just use the term \mathcal{C} -module. A \mathcal{C} -module homomorphism $\eta : M \rightarrow N$ between two \mathcal{C} -modules M and N is a natural transformation of functors. The category of all \mathcal{C} -modules forms a category, which we call $\mathbf{Rep}_{\mathbf{k}}(\mathcal{C})$.

A \mathcal{C} -module homomorphism $\eta : M \rightarrow N$ is *injective* (resp. *surjective*) if for each object $C \in \mathcal{C}$, the component $\eta_C : M(C) \rightarrow N(C)$ is injective (resp. surjective). We say N is a *submodule* (resp. *quotient module*) of M if there is an injective (resp. surjective) homomorphism $N \rightarrow M$ (resp. $M \rightarrow N$). It is a well-known fact that concepts such as subobjects, quotients, kernels, cokernels, images, direct sums, etc. can all be defined in this “pointwise” fashion in the context of \mathcal{C} -modules. In other words, $\mathbf{Rep}_{\mathbf{k}}(\mathcal{C})$ has the structure of an abelian category.

One of the key ingredients in [4] is the notion of a Noetherianity property. Recall that a module over a ring \mathbf{k} is Noetherian if every submodule is finitely generated (assuming the axiom of choice). The following definitions generalize these notions to \mathcal{C} -modules.

Definition 2.2 A \mathcal{C} -module M is *finitely generated* if there exist objects $C_1, C_2, \dots, C_n \in \mathcal{C}$ and elements $x_i \in M(C_i)$ for each i , satisfying that if N is a submodule of M such that $N(C_i)$ contains x_i , then $N = M$. The set $\{x_i\}$ is called the *generating set* of M .

Definition 2.3 A \mathcal{C} -module M is *Noetherian* if every submodule of M is finitely generated. The category of \mathcal{C} -modules is *locally Noetherian* if for any Noetherian ring \mathbf{k} , all finitely generated \mathcal{C} -modules are Noetherian.

An equivalent formulation of Definition 2.2 is occasionally useful. For any object $X \in \mathcal{C}$, let $P_{\mathcal{C}, X}$ denote the covariant representable \mathcal{C} -module generated at X , i.e. the functor defined by

$$\begin{aligned} P_{\mathcal{C}, X} : \mathcal{C} &\rightarrow \mathbf{Mod}_{\mathbf{k}} \\ Y &\rightarrow \mathbf{k}[\mathrm{Hom}_{\mathcal{C}}(X, Y)] \end{aligned}$$

for all $Y \in \mathcal{C}$. Then the following lemma holds:

Lemma 2.4 A \mathcal{C} -module is finitely generated if and only if it is a quotient of a direct sum of modules of the form $P_{\mathcal{C}, X}$.

Proof By the Yoneda lemma, a \mathcal{C} -module homomorphism $\eta : P_{\mathcal{C},X} \rightarrow M$ is determined uniquely by choosing an element $x \in M(X)$ and letting $\eta_X(1_X) = x$. It is straightforward to check that if M is finitely generated, then M is a quotient of the direct sum of the representable functors attached to the generating set. Similarly, if M is a quotient of this form, then the elements corresponding to each representable functor will be a generating set of M . \square

Let $f : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. This induces a functor $f^* : \mathbf{Rep}_k(\mathcal{D}) \rightarrow \mathbf{Rep}_k(\mathcal{C})$. The functor f is defined to be *finite* if for every $X \in \mathcal{D}$, $f^*(P_{\mathcal{D},X})$ is finitely generated. We end this subsection by recalling the following two lemmas, which appeared respectively as Lemmas 2.1 and 2.2 in [13].

Lemma 2.5 *Let \mathcal{C} be a category. The category of \mathcal{C} -modules is locally Noetherian if and only if for any object $X \in \mathcal{C}$, any submodule of $P_{\mathcal{C},X}$ is finitely generated.*

Lemma 2.6 *If the category of \mathcal{C} -modules is locally Noetherian, and $f : \mathcal{C} \rightarrow \mathcal{D}$ is a finite and essentially surjective functor, then the category of \mathcal{D} -modules is locally Noetherian.*

2.2 Semilocal Rings and Finite Rings

In this paper, we shall consider only finite commutative rings with unit. However, the literature of orthogonal forms often deal with semilocal rings, which are more general, so we briefly mention them here. Recall that a ring is *Artinian* if there is no infinite descending chain of ideals, and a ring R is *semilocal* if $R/\text{rad } R$ is Artinian. We recall an equivalent characterization of semilocal rings (c.f. [10]):

Proposition 2.7 *A ring R is semilocal if and only if it has finitely many maximal ideals.*

Therefore, it is clear that a finite ring is semilocal. Furthermore, if \mathfrak{p} is a prime ideal in a finite ring R , then R/\mathfrak{p} is a finite integral domain and therefore a field. This implies that \mathfrak{p} is maximal. Therefore, we have the following result (c.f. [10]):

Proposition 2.8 *A finite ring R is the direct product of finite local rings.*

Therefore, given a finite commutative ring R , we can express it as the product of finite local rings $R = \prod_{i=1}^n R_i$. Then, since each R_i is local, there is a unique maximal ideal \mathfrak{m}_i in each R_i , and this gives a projection map $\pi_i : R_i \rightarrow R_i/\mathfrak{m}_i$, the codomain of which is a field; in particular, the product map $\pi = \prod_{i=1}^n \pi_i$ gives a projection map from R to a product of finite fields R/\mathfrak{m} , where $\mathfrak{m} = \prod_{i=1}^n \mathfrak{m}_i$.

2.3 Symmetric Bilinear Forms

We now wish to define and characterize symmetric bilinear forms on finite commutative rings. As revealed shortly, we need to assume that 2 is a unit.

Definition 2.9 Let R be a semilocal ring, and let V be a finite-rank free R -module. A bilinear form $B : V \times V \rightarrow R$ is called *symmetric* or *orthogonal* if $B(v, w) = B(w, v)$ for all v, w . The form is said to be *non-degenerate* if it induces an isomorphism to the dual space $V^* = \text{Hom}_R(V, R)$. If B is non-degenerate, call the pair (V, B) an *orthogonal*

module. If (V, B_V) and (W, B_W) are two orthogonal modules, an R -module homomorphism $\phi : V \rightarrow W$ is called an *isometry* if $B_V(v, w) = B_W(\phi(v), \phi(w))$ for all $v, w \in V$.

From the definition, it follows that isometries are necessarily injective.

The classification of orthogonal modules over a finite ring up to bijective isometry is more difficult than the symplectic case. We first recall the following diagonalization theorem (c.f. [1]):

Proposition 2.10 *Let R be a semilocal ring, and let V be an orthogonal R -module. Then there exists a basis of V in which the matrix of B is diagonal and whose diagonal entries are units in R .*

In other words, we can find a bijective isometry from (V, B) to $(R^{\text{rk}V}, D)$, where D is a diagonal form as in the theorem. (Here, $\text{rk}V$ denotes the rank of a free R -module V .) While this theorem greatly simplifies the classification problem, it is still redundant (for instance, permuting basis vectors in $R^{\text{rk}V}$ will change D but the resulting module is still isometric). In the case where R is a finite field, the answer is well-known (though a proof is hard to find, c.f. [9]):

Proposition 2.11 *Let \mathbb{F} be a finite field (of characteristic $p > 2$), and let (V, B) be an orthogonal \mathbb{F} -module (i.e. a finite-dimensional vector space endowed with a non-degenerate symmetric bilinear form). Then, there exists a basis of V such that matrix of B is either 1) the identity matrix, or 2) the diagonal matrix $\text{diag}(1, \dots, 1, x)$, where x is any nonsquare in \mathbb{F}^\times , where different choices of x yield isometric forms.*

In other words, there are two isomorphism classes, and the dimension of V and the determinant of B determine the isomorphism class.

A similar result can be proven for a finite local ring. If R is a finite local ring, let $\pi : R \rightarrow \mathbb{F}$ denote the projection onto its residue field $\mathbb{F} = R/\mathfrak{m}$, where \mathfrak{m} is the maximal ideal in R . Then we have the following well-known proposition (we provide a proof because we could not find a proof in the literature):

Proposition 2.12 *Let R be a finite local ring (where 2 is a unit), and let (V, B) be an orthogonal R -module. Then, there exists a basis of V such that matrix of B is either 1) the identity matrix, or 2) the diagonal matrix $\text{diag}(1, \dots, 1, x)$, where $x \in R$ is such that $\pi(x)$ is a nonsquare in \mathbb{F}^\times , and where different choices of x yield isometric forms.*

Proof First of all, since R is a local ring, \mathfrak{m} consists of the non-units in R , so for any unit $u \in R$, the coset $u + \mathfrak{m}$ consists solely of units. By Proposition 2.10 and applying Proposition 2.11 to the induced orthogonal \mathbb{F} -module, we can find a basis $\{v_1, \dots, v_n\}$ of V such that with respect to this basis the form is diagonal, $B(v_i, v_j) = (1 + t_i)^{-1}$ where $t_i \in \mathfrak{m}$ for each $1 \leq i \leq n-1$ and B satisfies one of the following two cases: either $B(v_n, v_n) = (1 + t_n)^{-1}$ or $B(v_n, v_n) = (x + t_n)^{-1}$, where $t_n \in \mathfrak{m}$ and x is a unit in R such that $\pi(x)$ is a nonsquare in \mathbb{F}^\times .

Let us do case 1) first. For each i in $\{1, \dots, n\}$, consider the following quadratic equation in m : $(1 + m)^2 = 1 + t_i$, which can be rewritten as $m^2 + 2m - t_i = 0$. Since $t_i \in \mathfrak{m}$, reducing this monic polynomial modulo \mathfrak{m} gives a monic quadratic equation with two distinct roots $m(m + 2) = 0$, one of them being $m = 0$. By Theorem 3.12 in [8], it follows $m^2 + 2m -$

$t_i = 0$ has a root m_i in \mathfrak{m} . Then, in the basis $\{(1 + m_1)v_1, \dots, (1 + m_n)v_n\}$, we have $B((1 + m_i)v_i, (1 + m_i)v_i) = (1 + m_i)^2 B(v_i, v_i) = (1 + m_i)^2(1 + t_i)^{-1} = 1$ for all i .

For case 2), we can do the same thing for $1 \leq i \leq n - 1$. For $i = n$, we consider the polynomial equation $(x + m)^2 = x(x + t_n)$, which when reduced modulo \mathfrak{m} gives $m(m + 2\pi(x)) = 0$. The same reasoning gives a root $m = m_n \in \mathfrak{m}$ of $(x + m)^2 = x(x + t_n)$. Then, we have $B((x + m_n)v_n, (x + m_n)v_n) = (x + m_n)^2(x + t_n)^{-1} = x$. This proves the theorem. \square

Corollary 2.13 *Let R be a finite ring (where 2 is a unit), and write $R = \prod_{i=1}^q R_i$ as the product of finite local rings. Then, there are 2^q isomorphism classes of orthogonal R -modules.*

Proof Such a decomposition exists by Proposition 2.8. Let $e_i = (0, \dots, 0, 1_{R_i}, 0, \dots, 0)$ (nonzero in the i -th spot) be the central idempotent arising from $R_i = e_i R$, so $R = \bigoplus_{i=1}^n e_i R$. Then, since $1_R = e_1 + \dots + e_n$, it is clear that a bilinear form on R splits as the direct sum of bilinear forms on R_i . The result then follows from Theorem 2.12. \square

2.4 Column-Adapted Maps

In this last subsection, we discuss the notion of column-adapted and row-adapted maps, introduced in [13]. They are used in our proof of Lemma 3.4.

Definition 2.14 Let R be a commutative local ring. An R -linear map $f : R^m \rightarrow R^n$ is *column-adapted* if there is an n -element subset $S_c(f) = \{s_1 < s_2 < \dots < s_n\} \subseteq [m]$ such that, if we write f as a $n \times m$ matrix M with respect to the standard basis, then

- The s_i th column of M has 1 on the i th position and 0 elsewhere;
- The entries (i, j) where $j < s_i$ are all non-invertible.

For example, the map $f : R^5 \rightarrow R^3$ defined by the matrix

$$\begin{pmatrix} * & 1 & 0 & \bullet & 0 \\ * & 0 & 1 & \bullet & 0 \\ * & 0 & 0 & * & 1 \end{pmatrix}$$

is column adapted, if the entries labeled with $*$ are non-invertible (the entries labeled with \bullet can be any scalar).

In the general case where R is a finite commutative ring, by Proposition 2.8 there exists an isomorphism

$$R \cong R_1 \times \dots \times R_q$$

where the R_i 's are finite commutative local rings. In this case, we say a map $f : R^m \rightarrow R^n$ is *column-adapted* if the induced maps $R_i^m \rightarrow R_i^n$ are all column-adapted. Also, we say f is *row-adapted* if its transpose is column-adapted; in this case, we also define $S_r(f) = S_c(f^T)$.

The next two lemmas were established in [13] as Lemmas 2.9 and 2.10.

Lemma 2.15 *The composition of two column-adapted maps is column-adapted. Similarly, the composition of two row-adapted maps is row-adapted.*

Lemma 2.16 *Let R be a finite commutative ring, and let $f : R^{n'} \rightarrow R^n$ be a surjection. Then we can uniquely factor $f = f_2 f_1$, where $f_1 : R^{n'} \rightarrow R^n$ is column-adapted and $f_2 : R^n \rightarrow R^n$ is an isomorphism.*

3 Local Noetherianity

Throughout this section, R is a finite commutative ring where 2 is a unit, and \mathbf{k} is a Noetherian ring. Suppose we fix a factorization $R \cong R_1 \times R_2 \times \cdots \times R_q$, where each R_i is a local ring.

Definition 3.1 Define the following categories:

- $\mathbf{OrI}(R)$: objects are orthogonal R -modules (V, B) , and morphisms are isometries.
- $\mathbf{OrI}_{\text{sq}}(R)$: the full subcategory of $\mathbf{OrI}(R)$ spanned by objects (V, B) such that the projection of $\det B$ onto the residue fields of each R_i ($1 \leq i \leq q$) is a square.

We point out the important point that for any object $(V, B) \in \mathbf{OrI}(R)$, its group of automorphisms $\text{Aut}_{\mathbf{OrI}(R)}(V)$ is precisely the orthogonal group associated to (V, B) .

The goal in this section is to prove the following theorem:

Theorem 3.2 *The category $\mathbf{Rep}_{\mathbf{k}}(\mathbf{OrI}(R))$ is locally Noetherian.*

Our plan is to show the theorem with $\mathbf{OrI}(R)$ replaced with $\mathbf{OrI}_{\text{sq}}(R)$ in Section 3.1, and generalize it to the full category in Section 3.2.

3.1 The Square Case

In this subsection, our goal is to show that the category of $\mathbf{OrI}_{\text{sq}}(R)$ -modules is locally Noetherian (stated as Theorem 3.8 below). The proof is essentially identical to the analogous argument in [13].

Definition 3.3 Define the following categories:

- $\mathbf{OOrI}'_{\text{sq}}(R)$: objects are orthogonal R -modules (R^n, B) such that the projection from $\det B$ onto R_i is a square, and morphisms are row-adapted isometries.
- $\mathbf{OOrI}_{\text{sq}}(R)$: full subcategory of $\mathbf{OOrI}'_{\text{sq}}(R)$ spanned by objects which are orthogonal R -modules (R^n, B_{sq}) such that B_{sq} is the identity matrix under the standard basis.

Lemma 3.4 *Let R be a finite commutative ring where 2 is a unit. Let*

$$f \in \text{Hom}_{\mathbf{OrI}_{\text{sq}}(R)}((R^n, B), (R^{n'}, B')),$$

then we can uniquely write $f = f_1 f_2$ such that

- $f_2 \in \text{Hom}_{\mathbf{OrI}_{\text{sq}}(R)}((R^n, B), (R^n, \beta))$ where β has square determinant;
- $f_1 \in \text{Hom}'_{\mathbf{OOrI}_{\text{sq}}(R)}((R^n, \beta) \rightarrow (R^{n'}, B'))$.

Proof Transposing, applying Lemma 2.16, then transposing back, we can uniquely write $f = f_1 f_2$, where $f_1 : R^n \rightarrow R^{n'}$ is row-adapted and $f_2 : R^n \rightarrow R^n$ is an isomorphism. We can then uniquely choose β a symmetric form on R^n so that f_1 and f_2 are isometries. \square

Our first step in achieving the goal of this subsection is to show the existence of a well partial ordering \preceq on the set

$$\mathcal{P}_R(d, B) = \bigsqcup_{n \geq 0} \text{Hom}_{\mathbf{OOrI}_{\text{sq}}(R)}((R^d, B), (R^n, B_{\text{sq}}))$$

as described in the following lemma.

Lemma 3.5 *Fix R, d, B . There exists a well partial ordering \preceq on $\mathcal{P}_R(d, B)$ that can be extended to a total ordering \leq such that for $f, g \in \mathcal{P}_R(d, B)$, mapping to $R^n, R^{n'}$ respectively, satisfying $f \preceq g$, there exists some $\phi \in \text{Hom}_{\mathbf{OOrI}_{\text{sq}}(R)}((R^n, B_{\text{sq}}), (R^{n'}, B_{\text{sq}}))$ such that:*

- $g = \phi f$;
- For any $f_1 \in \text{Hom}_{\mathbf{OOrI}_{\text{sq}}(R)}((R^d, B), (R^n, B_{\text{sq}}))$ with $f_1 < f, \phi f_1 < g$.

To prove Lemma 3.5, we need to first show the following technical lemma. The proof of Lemma 3.5 is located shortly after it.

Lemma 3.6 *Fix R, d, B, f, g, n, n' as described in Lemma 3.5, except here we further restrict to the case where R is a finite commutative local ring. Let the rows of g be $r_1, \dots, r_{n'}$. Suppose that f can be obtained from g by deleting certain rows $r_{i_1}, \dots, r_{i_{n'-n}}$ where $I = \{i_1, \dots, i_{n'-n}\} \subseteq [n']$ such that $I \cap S_r(g) = \emptyset$. Then there exists $\phi \in \text{Hom}_{\mathbf{OOrI}_{\text{sq}}(R)}(R^n, R^{n'})$ such that:*

- For any $h \in \text{Hom}_{\mathbf{OOrI}_{\text{sq}}(R)}((R^d, B), (R^n, B_{\text{sq}}))$ with $S_r(h) = S_r(f)$, ϕh can be obtained from h by inserting the rows r_i in position i for each $i \in I$. In particular, $g = \phi f$.
- For any $h \in \text{Hom}_{\mathbf{OOrI}_{\text{sq}}(R)}((R^d, B), (R^n, B_{\text{sq}}))$ with $S_r(h) < S_r(f)$ in lexicographic order, then $S_r(\phi h) < S_r(g)$ in lexicographic order.

Proof The desired map ϕ can be defined by the following $n' \times n$ matrix: take a $(n' - n) \times n$ matrix whose k th row \widehat{r}_k is obtained from r_{i_k} by replacing the entries not in $S_r(f)$ with zeros. Then, we shuffle the rows of this matrix with the rows of a $n \times n$ identity matrix, such that the former rows occupy exactly the indices in I . It is straightforward to check that the two points hold and ϕ preserves the standard symmetric forms.

As a toy example, let $(d, n, n') = (3, 4, 6)$, and let

$$f = \begin{bmatrix} 1 & 0 & 0 \\ a & b & c \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } g = \begin{bmatrix} 1 & 0 & 0 \\ a & b & c \\ 0 & 1 & 0 \\ d & e & f \\ 0 & 0 & 1 \\ g & h & i \end{bmatrix}.$$

Then we take ϕ to be

$$\phi = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ d & 0 & e & f \\ 0 & 0 & 0 & 1 \\ g & 0 & h & i \end{bmatrix}.$$

Consider two vectors $v_i = (w_i, x_i, y_i, z_i)$ where $i \in \{1, 2\}$, let us now prove $B_{\text{sq}}(v_1, v_2) = B_{\text{sq}}(\phi(v_1), \phi(v_2))$. We know, by definition of f and g , that

$$\begin{aligned} B_{\text{sq}}(f(w_1, y_1, z_1), f(w_2, y_2, z_2)) &= B((w_1, y_1, z_1), (w_2, y_2, z_2)) \\ &= B_{\text{sq}}(g(w_1, y_1, z_1), g(w_2, y_2, z_2)), \end{aligned}$$

so

$$(dw_1 + ey_1 + fz_1)(dw_2 + ey_2 + fz_2) + (gw_1 + hy_1 + iz_1)(gw_2 + hy_2 + iz_2) = 0,$$

which is equivalent to $B_{\text{sq}}(v_1, v_2) = B_{\text{sq}}(\phi(v_1), \phi(v_2))$. In the general case, let the rows of f be f_i ($1 \leq i \leq n$), and the rows of g be g_i ($1 \leq i \leq n'$). Let $\alpha = (\alpha_i), \beta = (\beta_i)$ be any two vectors in R^n , we want to prove that $B_{\text{sq}}(\alpha, \beta) = B_{\text{sq}}(\phi(\alpha), \phi(\beta))$. Let α', β' denote the vectors in R^d obtained from α, β by only selecting the entries with indices in $S_r(f)$. Then

$$\begin{aligned} B_{\text{sq}}(\alpha, \beta) &= \sum_{i=1}^n \alpha_i \beta_i; \\ B_{\text{sq}}(\phi(\alpha), \phi(\beta)) &= \sum_{i=1}^n \alpha_i \beta_i + \sum_{i \in I} (g_i \cdot \alpha')(g_i \cdot \beta'); \\ B_{\text{sq}}(f(\alpha'), f(\beta')) &= \sum_{i=1}^n (f_i \cdot \alpha')(f_i \cdot \beta'); \\ B_{\text{sq}}(g(\alpha'), g(\beta')) &= \sum_{i=1}^{n'} (g_i \cdot \alpha')(g_i \cdot \beta'). \end{aligned}$$

Because $B_{\text{sq}}(f(\alpha'), f(\beta')) = B(\alpha, \beta) = B_{\text{sq}}(g(\alpha'), g(\beta'))$, we conclude that

$$B_{\text{sq}}(\alpha, \beta) = B_{\text{sq}}(\phi(\alpha), \phi(\beta)).$$

□

We now give the proof of Lemma 3.5.

Proof of Lemma 3.5 Assume first that R is a finite commutative local ring. Let $f, g \in \mathcal{P}_R(d, B)$ mapping to $R^n, R^{n'}$ respectively. We declare $f \leq g$ if f can be obtained from g by deleting some set of rows $I \subseteq [n']$ such that $I \cap S_r(g) = \emptyset$. This is clearly a partial order.

Let $\Sigma = R^d \sqcup \{\bullet\}$, \bullet being a formal symbol. Let Σ^* be the set of words whose letters come from Σ . There is a natural well-ordered poset structure on Σ^* (cf. [13] Lemma 2.5). We will prove that $(\mathcal{P}_R(d, B), \leq)$ is isomorphic to a subposet of Σ^* , which would imply that \leq is a well partial ordering. For $f \in \text{Hom}_{\mathbf{00r}_{\text{sq}}(R)}((R^d, B), (R^n, B_{\text{sq}}))$, let r_i represent the i th row of f if $i \notin S_r(f)$, else let $r_i = \bullet$. Thus, each $r_i \in \Sigma$, and we map f to the word $r_1 r_2 \dots r_n \in \Sigma^*$. Clearly this map is an order-preserving injection, so we conclude that \leq is a well partial ordering.

Next, we extend \leq to a total ordering \preceq . Fix an arbitrary total order on R^d . For $f \neq g$ the order is defined by

- If $n < n'$ then $f < g$;
- Otherwise, if $S_r(f) < S_r(g)$ in lexicographic order, then $f < g$;
- Otherwise, compare the sequences of rows of f and g by lexicographic order and the total order on R^d .

This clearly extends \leq , and the claimed properties follow by taking ϕ as described by Lemma 3.6. This shows Lemma 3.5 in the case where R is a finite commutative local ring.

In the general case where R is a finite commutative ring, fix an isomorphism

$$R \cong R_1 \times R_2 \times \dots \times R_q$$

where each R_i is a commutative local ring. Then

$$\text{Hom}_{\mathbf{OOrI}'_{\text{sq}}(R)}(R^n, R^{n'}) = \text{Hom}_{\mathbf{OOrI}'_{\text{sq}}(R_1)}(R_1^n, R_1^{n'}) \times \dots \times \text{Hom}_{\mathbf{OOrI}'_{\text{sq}}(R_q)}(R_q^n, R_q^{n'}).$$

This implies that each element $f \in \mathcal{P}_R(d, B)$ can be viewed as a tuple

$$(f_1, \dots, f_q) \in \mathcal{P}_{R_1}(d, B) \times \dots \times \mathcal{P}_{R_q}(d, B),$$

where each f_i maps into $R_i^{n'}$ for fixed n' . We can construct a partial order on the product by using the product partial order (where one element is smaller than another iff every component is). Then, extend this to a total order by lexicographic order. This restricts to a total order on $\mathcal{P}_R(d, B)$ that satisfies the necessary assumptions of Lemma 3.5. □

Using this well ordering, we can deduce the following theorem. The proof closely follows ideas in Section 4 of [14].

Theorem 3.7 *For $d \geq 0$ and B a symmetric form on R^d , any $\mathbf{OOrI}_{\text{sq}}(R)$ -submodule of the $\mathbf{OOrI}_{\text{sq}}(R)$ -module*

$$Q_{d,B} = \mathbf{k}[\text{Hom}_{\mathbf{OOrI}'_{\text{sq}}(R)}((R^d, B), -)]$$

is finitely generated. As a corollary, the category of $\mathbf{OOrI}_{\text{sq}}(R)$ -modules is locally Noetherian.

Proof In view of Lemma 2.5, it suffices to prove that any submodule of $Q_{d,B}$ is finitely generated (since we can take $B = B_{\text{sq}}$).

Fix d, B, R, \mathbf{k} , so we abbreviate $Q_{d,B}$ as Q . For an element $f \in \text{Hom}_{\mathbf{OOrI}'_{\text{sq}}(R)}((R^d, B), R^n)$, let e_f denote the basis vector in $Q(R^n)$ corresponding to f . For an element $x \in Q(R^n)$, define its **initial term** $\text{init}(x)$ as follows: if f is \leq -maximal such that e_f has coefficient $\alpha_f \neq 0$ in x , $\text{init}(x) = \alpha_f e_f$. Let M be a submodule of Q , we also define $\text{init}(M)$ to be a function taking R^n to the \mathbf{k} -module $\mathbf{k}[\text{init}(x) \mid x \in M(R^n)]$.

We claim that if N is a submodule of M and $N \neq M$, then $\text{init}(N) \neq \text{init}(M)$. Suppose for contradiction that $\text{init}(N) = \text{init}(M)$. Pick $y \in M(R^n) \setminus N(R^n)$ such that $\text{init}(y) = \alpha_t e_t$ is \leq -minimal. Since $\text{init}(M) = \text{init}(N)$, there exists $z \in N(R^n)$ such that $\text{init}(z) = \text{init}(y)$, but then $z - y \notin N(R^n)$ and $\text{init}(z - y)$ is smaller than e_t , contradiction. This proves the claim.

Suppose now that there exists an increasing sequence of submodules of Q

$$M_0 \subsetneq M_1 \subsetneq M_2 \subsetneq \dots$$

The claim implies that $\text{init}(M_i - 1) \neq \text{init}(M_i)$, so there exists, for every $i \geq 1$, some $n_i \geq 0$ and $\lambda_i e_{f_i} \in \text{init}(M_i)(R^{n_i}) \setminus \text{init}(M_{i-1})(R^{n_i})$. Because \preceq is a well partial ordering, there exists an infinite sequence $i_0 < i_1 < i_2 < \dots$ such that

$$f_{i_0} \preceq f_{i_1} \preceq f_{i_2} \preceq \dots$$

Since \mathbf{k} is Noetherian, we can choose m such that $\lambda_{i_m} = \sum_{j=0}^{m-1} c_j \lambda_{i_j}$ for $c_j \in \mathbf{k}$. For each $0 \leq j \leq m-1$, let $x_j \in M_{i_j}(R^{n_{i_j}})$ such that $\text{init}(x_j) = \lambda_{i_j} e_{f_{i_j}}$. By Lemma 3.5, there exists $\phi_j \in \text{Hom}_{\mathbf{OO}\mathbf{rI}_{\text{sq}}}(R^{i_j}, R^{i_m})$ such that $\phi_j f_{i_j} = f_{i_m}$ and for any $f'_{i_j} < f_{i_j}$ in the same Hom set, $\phi_j f'_{i_j} < f_{i_m}$.

Consider the element $X = \sum_{j=0}^{m-1} c_j \phi_j x_j$, which belongs to $M_{i_m-1}(R^{i_m})$. Then the properties in Lemma 3.5 implies that $\text{init}(X) = \lambda_{i_m} e_{f_{i_m}} \notin M_{i_m-1}(R^{i_m})$, contradiction. \square

Finally, we are ready to show that the category of $\mathbf{OrI}_{\text{sq}}(R)$ -modules is locally Noetherian.

Theorem 3.8 *The category of $\mathbf{OrI}_{\text{sq}}(R)$ -modules is locally Noetherian.*

Proof By Lemma 2.6 and Theorem 3.7, it suffices to show that the inclusion functor $\Phi : \mathbf{OO}\mathbf{rI}_{\text{sq}}(R) \rightarrow \mathbf{OrI}_{\text{sq}}(R)$ is finite (the essential surjectivity of Φ is obvious). Fix d, B , and let $M = P_{\mathbf{OrI}_{\text{sq}}(R), (R^d, B)}$, it suffices to prove that the $\mathbf{OO}\mathbf{rI}_{\text{sq}}(R)$ -module $\Phi^* = M \circ \Phi$ is finitely generated. Recall that by Theorem 3.7, $Q_{d, B'}$ is finitely generated for any symmetric form B' with square determinant on R^d . If we fix B' and an isometry $\tau : (R^d, B) \rightarrow (R^d, B')$, then we get a natural transformation $Q_{d, B'} \rightarrow \Phi^*$, and the map

$$\bigoplus_{B'} \bigoplus_{\tau} Q_{d, B'} \rightarrow \Phi^*(M)$$

is surjective by Lemma 3.4. It follows then that $\Phi^*(M) = M \circ \Phi$ is finitely generated, as desired. \square

3.2 The General Case

In the general case, we define the following subcategories of $\mathbf{OrI}(R)$: for each nonempty subset I of $\{1, 2, \dots, q\}$, let $\mathbf{OrI}_I(R)$ be the full subcategory of $\mathbf{OrI}(R)$ spanned by the objects (V, B) , where $\det B$ is a nonsquare in the residue field of R_i for all $i \in I$, and is a square otherwise. By Corollary 2.13, there are $2^q - 1$ of these categories. For each of them, we will show:

Theorem 3.9 *The category of $\mathbf{OrI}_I(R)$ -modules is locally Noetherian.*

Proof Define a functor $\Phi : \mathbf{OrI}_{\text{sq}}(R) \rightarrow \mathbf{OrI}_I(R)$, sending

$$(V, B) \mapsto (V, B) \oplus (R, X)$$

where X is the form on a 1-dimensional R -module sending 1 to $x = (x_1, x_2, \dots, x_q) \in R$, where x_i is an arbitrary nonsquare in the residue field of R_i if $i \in I$, and is 1 otherwise. This functor sends every isometry $f : (V, B) \rightarrow (W, B')$ in $\mathbf{OrI}_{\text{sq}}(R)$ to the isometry $\Phi(f) : (V, B) \oplus (R, X) \rightarrow (W, B') \oplus (R, X)$ where the last component is preserved. By

Lemma 2.6, because Φ is clearly essentially surjective, it suffices to check that it is finite, i.e. for any fixed $(V, B) \in \mathbf{OrI}_I(R)$, the $\mathbf{OrI}_{\text{sq}}(R)$ -module P mapping

$$(W, B') \mapsto \mathbf{k}[\text{Hom}_{\mathbf{OrI}_I(R)}((V, B), (W, B') \oplus (R, X))]$$

is finitely generated.

Consider the representable $\mathbf{OrI}_I(R)$ -module $P_{\mathbf{OrI}_I(R), (V, B)}$. This is clearly finitely generated. Pick a set of generators x_1, \dots, x_n . Then the same elements also generate P , which implies that it is finitely generated. \square

Using Theorems 3.8 and 3.9, we now give the proof of Theorem 3.2.

Proof of Theorem 3.2 Suppose that there exists a rising chain of $\mathbf{OrI}(R)$ -submodules

$$M^1 \subsetneq M^2 \subsetneq \dots \subset M$$

for some $\mathbf{OrI}(R)$ -module M . Restricting M and its submodules M^i to \mathbf{OrI}_{sq} , we get a chain of $\mathbf{OrI}_{\text{sq}}(R)$ -submodules

$$M_{\text{sq}}^1 \subsetneq M_{\text{sq}}^2 \subsetneq \dots \subset M_{\text{sq}}$$

Theorem 3.8 implies that this chain must stabilize at some finite N_{sq} . Similarly, restricting to each \mathbf{OrI}_I , we get a chain of $\mathbf{OrI}_I(R)$ -submodules

$$M_I^1 \subsetneq M_I^2 \subsetneq \dots \subset M_I$$

Theorem 3.9 implies that each such chain must stabilize at some finite N_I . Therefore, the original chain must also stabilize at a finite point, namely $1 + \max(N_{\text{sq}}, N_I)$. \square

4 Asymptotic Structure Theorem

4.1 Preliminaries

Let R be a finite commutative ring where 2 is a unit. Recall that a *complemented category* is a symmetric monoidal category (\mathcal{C}, \otimes) satisfying that:

- The identity object in \mathcal{C} is initial (hence there are *canonical morphisms* $V \rightarrow V \otimes V'$ and $V' \rightarrow V \otimes V'$);
- Every morphism in \mathcal{C} is a monomorphism;
- The map $\text{Hom}_{\mathcal{C}}(V \otimes V', W) \rightarrow \text{Hom}_{\mathcal{C}}(V, W) \times \text{Hom}_{\mathcal{C}}(V', W)$, defined by composing with the canonical morphisms, is injective;
- For every subobject V of W , there exists a unique subobject V' of W such that there is an isomorphism $V \otimes V' \rightarrow W$, which satisfies that the compositions $V \rightarrow V \otimes V' \rightarrow W$ and $V' \rightarrow V \otimes V' \rightarrow W$ are the inclusions.
- The map $S_n \curvearrowright \text{Aut}_{\mathcal{C}}(V) \rightarrow \text{Aut}_{\mathcal{C}}(V^n)$ is injective.

Furthermore, a complemented category \mathcal{C} is *generated* by an object X if any object is isomorphic to X^n for a unique $n \geq 0$. We recall the following theorem, whose three-part structure reflects a parallel with the multiplicity stability theorem described in Section 1:

Theorem 4.1 (Theorem F, [13]) *Let (\mathcal{C}, \otimes) be a complemented category with a generator X ; assume that the category of \mathcal{C} -modules is locally Noetherian, and let M be a finitely generated \mathcal{C} -module. For $N \geq 0$, denote by \mathcal{C}^N the full subcategory of \mathcal{C} generated by all objects of X -rank at most N . Then:*

- (Injective stability) If $f : V \rightarrow W$ is a morphism in \mathcal{C} , then $M(f) : M(V) \rightarrow M(W)$ is injective when the X -rank of V is sufficiently large.
- (Surjective stability) If $f : V \rightarrow W$ is a morphism in \mathcal{C} , then the orbit under $\text{Aut}_{\mathcal{C}}(W)$ of the image of $M(f)$ spans $M(W)$ when the X -rank of V is sufficiently large.
- (Central stability) For N sufficiently large, the functor M is the left Kan extension to \mathcal{C} of the restriction of M to \mathcal{C}^N .

Because the category $\text{OrI}_{\text{sq}}(R)$ is a complemented category generated by the 1-dimensional R -module $(R, 1)$, and the category of $\text{OrI}_{\text{sq}}(R)$ -modules is locally Noetherian by Theorem 3.8, Theorem 4.1 holds for $\text{OrI}_{\text{sq}}(R)$. In this section, we will prove an analogous theorem for $\text{OrI}(R)$ (Theorems 4.5 and 4.13 and Corollary 4.9).

4.2 Surjective Stability

For simplicity, we will use \mathcal{A} in place for $\text{OrI}(R)$ in this section.

Lemma 4.2 *The category $\mathcal{A} = \text{OrI}(R)$ is a complemented category.*

Proof The monoidal structure is given by the orthogonal direct sum, and for a free orthogonal submodule $V \subseteq W$ the complement of V is given by $V^\perp = \{w \in W \mid B(v, w) = 0\}$, where B is the nondegenerate symmetric form equipped on W . □

We remark that \mathcal{A} is generated by not one generator, but instead 2^q , one for each subset $I \subseteq \{1, 2, \dots, q\}$: we denote by \mathcal{X}_I the 1-dimensional free R module equipped with the bilinear form $x = (x_1, \dots, x_q)$ where x_i is a fixed nonsquare in the residue field of R_i if $i \in I$, and $x_i = 1$ otherwise. The fact that they generate \mathcal{A} follows from Proposition 2.13. We also remark that \mathcal{X}_I^2 is isomorphic to $\mathcal{X}_{\emptyset}^2$, because in the residue fields of R_i , the product of two nonsquares is a square.

Lemma 4.3 *For $V, W \in \mathcal{A}$, $\text{Aut}_{\mathcal{A}}(W)$ acts transitively on $\text{Hom}_{\mathcal{A}}(V, W)$.*

Proof Suppose we are given morphisms $f, g \in \text{Hom}_{\mathcal{A}}(V, W)$. Suppose the complements of $f(V)$ and $g(V)$ in W are respectively U, U' . The orthogonal modules $f(V), g(V)$ are both isomorphic to V , and the orthogonal modules U, U' are isomorphic as well. So we get the automorphism

$$\phi : W \cong f(V) \oplus U \xrightarrow{\cong} g(V) \oplus U' \cong W$$

satisfies $\phi f = g$, as desired. □

Corollary 4.4 *Let $f : U \rightarrow V$ and $g : U \rightarrow W$ be morphisms in \mathcal{A} , such that $\text{rk}V < \text{rk}W$. Then there exist a morphism $h : V \rightarrow W$ such that $h \circ f = g$.*

Proof Because $\text{rk}V < \text{rk}W$, we claim there exists a morphism $i : V \rightarrow W$. To see this, suppose B, B' are the bilinear forms attached to V and W , and $\det B$ is nonsquare in R_i for $i \in I_V$ while $\det B'$ is nonsquare in R_i for $i \in I_W$. Then $W \cong V \oplus (R, \mathcal{X}_{I_V \Delta I_W})$ (Δ means set XOR).

By Lemma 4.3, there exists an automorphism $k : W \rightarrow W$ such that $k \circ g = i \circ f$, hence $g = (k^{-1} \circ i) \circ f$. □

Theorem 4.5 (Surjective stability) *Let M be a finitely generated \mathcal{A} -module. Then surjective stability holds: for an \mathcal{A} -morphism $f : V \rightarrow W$ with $\text{rk}V$ large enough, the image of $M(f)$ spans $M(W)$ under $M(\text{Aut}_{\mathcal{A}}(W))$.*

Proof Suppose $f : V \rightarrow W$ is any morphism in \mathcal{A} . The span of $M(f)$ in $M(W)$, under the action of $M(\text{Aut}_{\mathcal{A}}(W))$, is the span of $M(\phi \circ f)$ as ϕ ranges in $\text{Aut}_{\mathcal{A}}(W)$. By Lemma 4.3, this is the same as the span of $M(h)$ as h ranges in $\text{Hom}_{\mathcal{A}}(V, W)$.

Now, because M is finitely generated, there exists a generating set $\{x_i \in M(V_i)\}$. Let $r = \max_i \text{rk}V_i$, and consider any object V with $\text{rk}V \geq r + 1$. Fix maps $\phi_i : V_i \rightarrow V$. For any $x \in W$, since M is generated by the x_i 's, there exist maps $f_i : V_i \rightarrow W$ such that x is in the span of the images of $M(f_i) : M(V_i) \rightarrow M(W)$.

If $\text{rk}V < \text{rk}W$, then by Corollary 4.4 there exist maps $h_i : V \rightarrow W$ such that $h_i \circ \phi_i = f_i$. This implies that x lies in the span of $M(h_i)$, as desired. If $\text{rk}V = \text{rk}W$, then since there exists a map $f : V \rightarrow W$, V must be isomorphic to W , so the proof of Corollary 4.4 still applies, and x lies in the span of $M(h_i)$. □

4.3 Injective and Central Stability

We will now prove injective and central stability.

Definition 4.6 Let M be an \mathcal{A} -module. The *torsion submodule* M_T of M is defined by

$$M_T(V) = \{x \in M(V) \mid \exists f : V \rightarrow W, M(f)(x) = 0\}.$$

Lemma 4.7 *The torsion submodule M_T is an \mathcal{A} -submodule of M .*

Proof It suffices to show that if $x, y \in M_T(V)$, then $x + y \in M_T(V)$. Suppose $f : V \rightarrow W$ and $g : V \rightarrow W'$ such that $M(f)(x) = M(g)(y) = 0$. Consider the maps $\iota : W \hookrightarrow W \oplus W'$ and $\iota' : W' \hookrightarrow W \oplus W'$. By Lemma 4.3, there exists $h \in \text{Aut}_{\mathcal{A}}(W \oplus W')$ such that $h\iota'g = \iota f$. This composition maps both x and y to 0, hence also $x + y$. □

Lemma 4.8 *Let M be finitely generated, then for all $V \in \mathcal{A}$ with $\text{rk}V \gg 0$, $M_T(V) = 0$.*

Proof Because the category of \mathcal{A} -modules is locally Noetherian and M_T is a submodule of M , M_T is finitely generated as well. Then all maps into sufficiently large-rank spaces are zero maps by Corollary 4.4. □

Corollary 4.9 (Injective stability) *Let M be a finitely generated \mathcal{A} -module. Then injective stability holds: for an \mathcal{A} -morphism $f : V \rightarrow W$ with $\text{rk}V$ large enough, $M(f)$ is injective.*

Proof This is a direct consequence of the above lemma. □

Definition 4.10 Let n be a positive integer, and let M be an \mathcal{A} -module. Suppose I is a subset of $\{1, 2, \dots, q\}$. Define $\Sigma_{I,n}M$ to be an \mathcal{A} -module mapping each $V \in \mathcal{A}$ to

$$\Sigma_{I,n}M(V) = \bigoplus_{h \in \text{Hom}_{\mathcal{A}}(\mathcal{X}_I^n, V)} M(C_h),$$

where C_h denotes the complement of $h(\mathcal{X}_I^n)$ in V .

Lemma 4.11 *If M is finitely generated, then the \mathcal{A} -modules $\Sigma_{I,n}M$ are all finitely generated.*

Proof Because M is finitely generated, there exists a set of generators x_1, \dots, x_m , which respectively belong to $M(V_1), \dots, M(V_m)$. Notice that each $M(V_i)$ is a direct summand in $\Sigma_{I,n}M(V_i \oplus \mathcal{X}_I^n)$. We claim that the images of these elements x_1, \dots, x_m also generate $\Sigma_{I,n}M$. For any nonzero $\Sigma_{I,n}M(V)$, V must be isomorphic to $\mathcal{X}_I^n \oplus Y$ for some orthogonal module Y . Clearly, the direct summand $M(Y)$ inside $\Sigma_{I,n}M(\mathcal{X}_I^n \oplus Y)$ is generated by the claimed set of generators; by Lemma 4.3, the other direct summands are all generated by them as well. \square

We can endow $\Sigma_{I,\bullet}M$ with a chain complex structure as follows. Consider an object $V \in \mathcal{A}$, and we define the maps $d_1, \dots, d_n : \Sigma_{I,n}M(V) \rightarrow \Sigma_{I,n-1}M(V)$ as induced by the maps $\mathcal{X}_I^{n-1} \rightarrow \mathcal{X}_I^n$ by “adding” the i th coordinate ($1 \leq i \leq n$). Finally, define $d = d_1 - d_2 + \dots + (-1)^{n-1}d_n$. It is straightforward to check that $\Sigma_{\bullet}M$ is a chain complex.

Theorem 4.12 *Let M be finitely generated. Fix n a positive integer, and $I \subseteq \{1, 2, \dots, q\}$. Then the chain complex*

$$\Sigma_{I,n}M(V) \rightarrow \dots \rightarrow \Sigma_{I,1}M(V) \rightarrow M(V) \rightarrow 0$$

is exact for all V with sufficiently large rank (as free R -modules).

Proof It suffices to prove that $(\mathcal{H}_i(\Sigma_{I,\bullet}M))(V) = (\mathcal{H}_i(\Sigma_{I,\bullet}M)_T)(V)$, since then it follows from Lemma 4.8 that $(\mathcal{H}_i(\Sigma_{\bullet}M))(V) = 0$ for sufficiently large $\text{rk}V$, which implies the exactness of the chain complex. To prove the claimed fact, we only need to show that the map $(\mathcal{H}_i(\Sigma_{I,\bullet}M))(V) \rightarrow (\mathcal{H}_i(\Sigma_{I,\bullet}M))(V \oplus \mathcal{X}_I)$, induced by the canonical morphism $V \rightarrow V \oplus \mathcal{X}_I$, is the zero map. But then the argument in Lemma 3.11 of [13] applies verbatim. \square

Now, we are ready to show central stability. The argument closely follows the proof of Theorem F in [13]. The main modification is the use of the chain complex $\Sigma_{I,n}M$, which is parametrized by I to handle the various isomorphism classes of orthogonal modules for any given rank.

Theorem 4.13 (Central stability) *Let $\mathcal{A}_{\leq N}$ denote the full subcategory spanned by the objects with R -rank at most N . Let M be a finitely generated \mathcal{A} -module. Then for all N sufficiently large, M is the left Kan extension to $M|_{\mathcal{A}_{\leq N}}$ along the inclusion functor $p : \mathcal{A}_{\leq N} \rightarrow \mathcal{A}$.*

Proof In the proof that follows, define $\mathcal{A}_{\leq N}$ instead to be the full subcategory spanned by objects of the following form:

- objects with R -rank at most $N - 1$;
- objects isometric to the rank- N orthogonal R -module equipped with the bilinear form whose matrix representation under the standard basis is the identity.

The reason for doing this is because of a minor problem in factoring morphisms. This does not change what we needed to show, since restricting to a larger set of objects does not change the left Kan extension (as can be easily seen from its universal property).

Let M' be the desired left Kan extension, then the universal property gives a natural transformation $\phi : M' \rightarrow M$ such that $\phi_V : M'(V) \rightarrow M(V)$ are isomorphisms for all V with $\text{rk} V \leq N - 1$. (Here we choose N sufficiently large so that $\Sigma_{I,2}M \rightarrow \Sigma_{I,1}M \rightarrow M \rightarrow 0$ is exact whenever $\text{rk} V \geq N$.) It suffices to prove that ϕ_V are isomorphisms for all V . Unlike in the proof of Theorem F in [13], the chain complex $\Sigma_{I,n}M$ we choose will depend on the isomorphism class of V , which is the key difference from that proof.

Induct on $\text{rk} V$, and we assume ϕ_V is an isomorphism for all $\text{rk} V \leq r - 1$ ($r \geq N + 1$). Fix V to be a rank- r object, and without loss of generality $V = \mathcal{X}_\emptyset^{r-1} \oplus \mathcal{X}_{I_0}$. The natural transformation ϕ induces natural transformations $\Sigma_{I,i}M \rightarrow \Sigma_{I,i}M'$ for each I , and by definition we know the induced maps $\Sigma_{I,i}M(V) \rightarrow \Sigma_{I,i}M'(V)$ are isomorphisms for each $i \geq 1$ and $I \subseteq \{1, \dots, q\}$.

Consider the commutative diagram

$$\begin{array}{ccccccc}
 \Sigma_{I_0,2}M'(V) & \longrightarrow & \Sigma_{I_0,1}M'(V) & \longrightarrow & M'(V) & \longrightarrow & 0 \\
 \downarrow \cong & & \downarrow \cong & & \downarrow \phi_V & & \\
 \Sigma_{I_0,2}M(V) & \longrightarrow & \Sigma_{I_0,1}M(V) & \longrightarrow & M(V) & \longrightarrow & 0,
 \end{array} \tag{1}$$

whose bottom row is exact. We now wish to prove that the map $\Sigma_{I_0,1}M'(V) \rightarrow M'(V)$ is surjective, from which it will follow that ϕ_V is an isomorphism by a simple diagram chase on (1).

By definition of the Kan extension, for $V \in \mathcal{A}$, $M'(V)$ is the colimit

$$\varinjlim((p \downarrow V) \rightarrow \mathcal{A}_{\leq N} \xrightarrow{M} \mathbf{Mod}_{\mathbf{k}}).$$

Furthermore, it is easy to see that is a filtered colimit (due to our definition of $\mathcal{A}_{\leq N}$), so as a set it is explicitly given by

$$M'(V) = \left(\coprod_{\substack{(U,f):U \xrightarrow{f} V \\ U \in \mathcal{A}_{\leq N}}} M(U) \right) / \sim$$

where \sim is the usual equivalence relation. By Corollary 4.4, any map $f : U \rightarrow V$ from an object U of rank at most N to V factors through an object W of rank $r - 1$. Furthermore, we can suppose $W = C_h$, where $h : \mathcal{X}_{I_0} \rightarrow V$ and C_h is the complement of the image of h in V . The inclusion $W \rightarrow V$ induces a map $u : M'(W) \rightarrow M'(V)$. Define another map $u' : M'(W) \rightarrow M'(V)$ by the universal property of $M'(W)$ as a filtered colimit. It suffices to show that u and u' are the same map, since then it would imply the surjectivity of $\Sigma_{I_0,1}M'(V) \rightarrow M'(V)$.

To do this, consider the objects $\{M'(U) : U \in \mathcal{A}_{\leq N}, f : U \rightarrow V\}$. These, along with the morphisms $M'(U) \rightarrow M'(U')$ defined by

$$M'(U) \xrightarrow{\cong} M(U) \rightarrow M(U') \xrightarrow{\cong} M'(U'),$$

form another diagram of shape $(p \downarrow W)$ in \mathcal{A} . The colimit of this is again $M'(W)$, and the map $u : M'(W) \rightarrow M'(V)$ is precisely the (unique) map provided by the universal property. Therefore, we conclude that $u = u'$, which finishes the proof. □

Remark 4.14 We would like to present an alternative proof of central stability, pointed to us by a referee of the paper, using a condition for central stability proven in [7]. Namely, given a category \mathcal{A} whose set of objects is $\mathbb{Z}_{\geq 0}$, an \mathcal{A} -module \mathcal{M} can be regarded as a graded module M over the category algebra A of \mathcal{A} (see these definitions in section 2 of [7]). Let $e_n \in A$ be the idempotent representing the identity morphism on the object $n \in \text{Ob}(\mathcal{A})$, and define

$$e_{0,N} = e_0 + e_1 + \dots + e_N.$$

We call M *centrally stable* if for all N large enough,

$$Ae \otimes_{eAe} eM \cong M,$$

where $e = e_{0,N}$. Then Proposition 2.4 in [7] shows that M is centrally stable if and only if \mathcal{M} satisfies central stability in the sense of Theorem F, [13]. With little modifications, the same can be shown to be true for **OrI**(R)-modules: take \mathcal{A} to be a skeleton of **OrI**(R), and $e_{0,N}$ to be the sum of elements representing the identity morphisms of all objects in \mathcal{A} of rank at most N , then an **OrI**(R)-module (equivalently, an \mathcal{A} -module) \mathcal{M} is centrally stable in the sense of Theorem 4.13 if and only if the associated A -module M is centrally stable.

This gives a simpler proof of Theorem 4.13: suppose that \mathcal{M} is a finitely generated \mathcal{A} -module, then M is finitely presented over $Ae_{0,N}$ for all sufficiently large N , by Theorem 3.2 (local Noetherianity). Then Lemma 3.1 in [7] shows that $Ae \otimes_{eAe} eM \cong M$ for $e = e_{0,N}$ for N sufficiently large, so M is indeed centrally stable, which implies that \mathcal{M} satisfies central stability.

5 Twisted and Untwisted Homological Stability

In this section, we fix R to be a finite field with $\text{char } R > 2$. In this case, there are 2 isomorphism classes of non-degenerate symmetric bilinear forms in each dimension.

Define, for an object $(V, B) \in \mathbf{OrI}(R)$, $O(V, B)$ to be the orthogonal group associated with (V, B) (i.e. the group of R -linear isomorphisms $V \rightarrow V$ preserving the bilinear form B). Then for any morphism $(V, B) \rightarrow (W, B')$, there is an induced map $O(V, B) \rightarrow O(W, B')$ given by mapping f to $f \oplus \text{id}$, where id is the identity map on the complement of (V, B) in (W, B') .

In this section, we prove the following twisted homological stability theorem:

Theorem 5.1 *Let R be a finite field with $\text{char } R > 2$. Let M be a finitely generated **OrI**(R)-module over $\mathbf{k} = \mathbb{Z}/\ell\mathbb{Z}$, where ℓ is a prime not equal to $\text{char } R$. Fix any index $k \geq 0$. Consider a morphism $(V, B) \rightarrow (W, B')$ in **OrI**(R). As explained above, this induces a map on the homologies*

$$H_k(O(V, B); M(V, B)) \rightarrow H_k(O(W, B'); M(W, B')).$$

Then this map is an isomorphism for all V with sufficiently large rank n .

First, we need to state several lemmas.

Lemma 5.2 *Let $V, V' \in \mathbf{OrI}(R)$, and let $i : V \rightarrow V \oplus V'$ be the canonical injection given by $v \mapsto v \oplus 0$. Then*

$$\{\tau \in \text{Aut}(V \oplus V') : \tau i = i\} \cong \text{Aut}(V').$$

Proof See the proof of [13], Lemma 3.2. Because $\mathbf{OrI}(R)$ is a complemented category, the conclusion there applies to $\mathbf{OrI}(R)$. \square

Lemma 5.3 *Let $V, W \in \mathbf{OrI}(R)$, then, as $\text{Aut}(V)$ -representations,*

$$\mathbf{k}[\text{Hom}(V, V \oplus W)] \cong \mathbf{k}[\text{Aut}(V \oplus W)/\text{Aut}(W)] \cong \text{Ind}_{\text{Aut}(W)}^{\text{Aut}(V \oplus W)}(\mathbf{k}).$$

Proof By Lemma 4.3, $\text{Aut}(V \oplus W)$ acts transitively on $\text{Hom}(V, V \oplus W)$. By Lemma 5.2, the stabilizer of any element $i \in \text{Hom}(V, V \oplus W)$ is isomorphic to $\text{Aut}(W)$, which proves the first equality. The second equality follows by definition of an induced representation. \square

Corollary 5.4 *For any $V, W \in \mathbf{OrI}(R)$,*

$$H_k(\text{Aut}(V \oplus W); \mathbf{k}[\text{Hom}(V, V \oplus W)]) \cong H_k(\text{Aut}(W); \mathbf{k}).$$

Proof We have

$$\begin{aligned} H_k(\text{Aut}(W); \mathbf{k}) &\cong H_k(\text{Aut}(V \oplus W); \text{Ind}_{\text{Aut}(W)}^{\text{Aut}(V \oplus W)}(\mathbf{k})) \\ &\cong H_k(\text{Aut}(V \oplus W); \mathbf{k}[\text{Hom}(V, V \oplus W)]). \end{aligned}$$

where the first step uses Shapiro’s lemma for group homology ($\text{Aut}(W) \leq \text{Aut}(V \oplus W)$ by acting trivially on V), and the second step uses Lemma 5.3). \square

Now, we prove the main theorem of this section.

Proof of Theorem 5.1 Let X, Y respectively denote the square and nonsquare generator of $\mathbf{OrI}(R)$; in the notation of Section 4, they would be $X = \mathcal{X}_\emptyset$ and $Y = \mathcal{X}_{\{1\}}$. Also, fix an isomorphism $X^2 \cong Y^2$

First, we consider the case where we impose the following two extra conditions on M :

- (i) For any morphism $f : V \rightarrow W$ in $\mathbf{OrI}(R)$, the map $M(f) : M(V) \rightarrow M(W)$ is injective;
- (ii) There exists a set of generators of M such that each generator lies in $M(V)$, where $V \cong X^r \oplus Y$ for some $r \geq 0$ (possibly depending on the generator).

In this case, we will show that for any fixed k , the map

$$H_k(\text{Aut}(X^n \oplus Y); M(X^n \oplus Y)) \rightarrow H_k(\text{Aut}(X^{n+1} \oplus Y); M(X^{n+1} \oplus Y)) \tag{2}$$

is an isomorphism for all $n \gg 0$.

Because M is finitely generated, by Lemma 2.4, M is a quotient module of a projective module P_0 , which is the direct sum of finitely many representable $\mathbf{OrI}(R)$ -modules. Furthermore, because of condition (ii), we can choose P_0 so that it is the direct sum of representable functors based at objects isomorphic to those of the form $X^r \oplus Y$. The kernel of $P_0 \rightarrow M$ is finitely generated by local Noetherianity. Consider the following commutative diagram, where $f : V \rightarrow W$ is a morphism in $\mathbf{OrI}(R)$:

$$\begin{array}{ccc} P_0(V) & \xrightarrow{P_0(f)} & P_0(W) \\ \downarrow & & \downarrow \\ M(V) & \xrightarrow{M(f)} & M(W). \end{array}$$

Because $M(f)$ is injective, we see that if $v \in P_0(V)$ and $P_0(f)(v) \in \ker(P_0(W) \rightarrow M(W))$, then $v \in \ker(P_0(V) \rightarrow M(V))$. From this, we deduce that the kernel $\ker(P_0 \rightarrow M)$ must also satisfy its analogous properties (i) and (ii).

Therefore, we can repeat this process to extend this to a projective resolution \bar{C} of M by direct sums of finitely many representable **OrI**(R)-modules based at objects isomorphic to $X^r \oplus Y$:

$$\bar{C} : \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0.$$

Now, if we delete M from the chain complex \bar{C} to produce

$$C : \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow 0,$$

then for V an object in **OrI**(R),

$$H_k(\text{Aut}(V); M(V)) \cong H_k(\text{Aut}(V); C),$$

where the right hand side is group homology with coefficients in a chain complex. There exists a spectral sequence (cf. [2]):

$$E_{pq}^1 = H_p(\text{Aut}(V); P_q(V)) \implies H_{p+q}(\text{Aut}(V); C). \tag{3}$$

Because each P_q is the direct sum of representable modules based at objects isomorphic to some $X^r \oplus Y$, we see (using Corollary 5.4) that for any fixed p, q, n , there is a map

$$\begin{aligned} \bigoplus_i H_p(\text{Aut}(X^i); \mathbf{k}) &\cong H_p(\text{Aut}(X^n \oplus Y); P_q(X^n \oplus Y)) \\ &\rightarrow H_p(\text{Aut}(X^{n+1} \oplus Y); P_q(X^{n+1} \oplus Y)) \cong \bigoplus_i H_p(\text{Aut}(X^{1+i}); \mathbf{k}). \end{aligned}$$

From [6], each map $H_p(\text{Aut}(X^i); \mathbf{k}) \rightarrow H_p(\text{Aut}(X^{1+i}); \mathbf{k})$ is an isomorphism for all $i \gg 0$. Hence, we conclude that

$$H_p(\text{Aut}(X^n \oplus Y); P_q(X^n \oplus Y)) \xrightarrow{\cong} H_p(\text{Aut}(X^{n+1} \oplus Y); P_q(X^{n+1} \oplus Y))$$

for all $n \gg 0$. Applying the spectral sequence (3), we conclude that for any fixed $k = p + q$, the map (2) is indeed an isomorphism for all $n \gg 0$.

Next, we consider the second case where M satisfies (i) and the following condition (ii'), instead of (ii):

- (ii') There exists a set of generators of M such that each generator lies in $M(V)$, where $V \cong X^r$ for some $r \geq 0$ (possibly depending on the generator).

Using the exact same argument, we can similarly show that for any fixed k , the map

$$H_k(\text{Aut}(X^n); M(X^n)) \rightarrow H_k(\text{Aut}(X^{n+1}); M(X^{n+1}))$$

is an isomorphism for all $n \gg 0$.

We now consider the general case where M is any finitely generated **OrI**(R)-module. Fix a large enough positive integer N according to Theorem 4.5 and Corollary 4.9, such that injective and surjective stability holds for all objects with rank at least N . Let M_1 be the submodule of M given by

$$M_1(V) = \begin{cases} M(V) & \text{if either } \text{rk} V > N \text{ or } V \cong X^{N-1} \oplus Y; \\ 0 & \text{otherwise,} \end{cases}$$

and for any morphism $f : V \rightarrow W$, $M_1(f)$ is the zero map when $M_1(V) = 0$, and is the same as $M(f)$ otherwise. Then by local Noetherianity, M_1 is a finitely generated module

satisfying conditions (i) and (ii), and $M_1(V) \cong M(V)$ for all V of large enough rank. As a result,

$$\begin{aligned} H_k(\text{Aut}(X^n \oplus Y); M(X^n \oplus Y)) &= H_k(\text{Aut}(X^n \oplus Y); M_1(X^n \oplus Y)) \\ &\cong H_k(\text{Aut}(X^{n+1} \oplus Y); M_1(X^{n+1} \oplus Y)) \\ &= H_k(\text{Aut}(X^{n+1} \oplus Y); M(X^{n+1} \oplus Y)) \end{aligned}$$

is an isomorphism for all $n \gg 0$.

Similarly, let M_2 be the submodule of M given by

$$M_2(V) = \begin{cases} M(V) & \text{if either } \text{rk} V > N \text{ or } V \cong X^N; \\ 0 & \text{otherwise,} \end{cases}$$

and for any morphism $f : V \rightarrow W$, $M_2(f)$ is the zero map when $M_2(V) = 0$, and is the same as $M(f)$ otherwise. Then by local Noetherianity, M_2 is a finitely generated module satisfying conditions (i) and (ii'), and $M_2(V) \cong M(V)$ for all V of large enough rank. As a result,

$$\begin{aligned} H_k(\text{Aut}(X^n); M(X^n)) &= H_k(\text{Aut}(X^n); M_2(X^n)) \\ &\cong H_k(\text{Aut}(X^{n+1}); M_2(X^{n+1})) \\ &= H_k(\text{Aut}(X^{n+1}); M(X^{n+1})) \end{aligned}$$

is an isomorphism for all $n \gg 0$.

Finally, because the composition of isometries $X^{n-2} \oplus Y \rightarrow X^n \rightarrow X^n \oplus Y \rightarrow X^{n+2}$, given by the inclusions and isomorphisms

$$X^{n-2} \oplus Y \rightarrow X^{n-2} \oplus Y^2 \cong X^n \rightarrow X^n \oplus Y \rightarrow X^n \oplus Y^2 \cong X^{n+2}$$

induces the maps

$$\text{Aut}(X^{n-2} \oplus Y) \rightarrow \text{Aut}(X^n) \rightarrow \text{Aut}(X^n \oplus Y) \rightarrow \text{Aut}(X^{n+2}),$$

which induces the maps

$$\begin{aligned} H_k(\text{Aut}(X^{n-2} \oplus Y); M(X^{n-2} \oplus Y)) &\xrightarrow{f} H_k(\text{Aut}(X^n); M(X^n)) \\ &\xrightarrow{g} H_k(\text{Aut}(X^n \oplus Y); M(X^n \oplus Y)) \\ &\xrightarrow{h} H_k(\text{Aut}(X^{n+2}); M(X^{n+2})), \end{aligned}$$

and since gf, hg are both isomorphisms, we conclude that g is an isomorphism as well. Because any object $(V, B) \in \mathbf{OrI}(R)$ is isomorphic to either X^n or $X^n \oplus Y$ ($n \geq 0$), these are enough to imply Theorem 5.1. \square

As a corollary, we show the following homological stability result with untwisted coefficients:

Corollary 5.5 *Under assumptions stated in Theorem 5.1, the map*

$$H_k(O(V, B); \mathbf{k}) \rightarrow H_k(O(W, B'); \mathbf{k})$$

is an isomorphism for all V with sufficiently large rank.

Proof Take m sufficiently large. Consider the finitely generated module $P_{\mathbf{OrI}(R), X^m}$. Then for any object $V \in \mathbf{OrI}(R)$,

$$\begin{aligned} H_k(\mathrm{Aut}(V); \mathbf{k}) &= H_k(\mathrm{Aut}(V \oplus X^m); \mathbf{k}[\mathrm{Hom}(X^m, V \oplus X^m)]) \\ &\cong H_k(\mathrm{Aut}(A \oplus X^{m+1}); \mathbf{k}[\mathrm{Hom}(X^m, V \oplus X^{m+1})]) \\ &= H_k(\mathrm{Aut}(V \oplus X); \mathbf{k}), \end{aligned}$$

where the first and third steps follow from 5.4, and the second step follows from Theorem 5.1 (recall we took m sufficiently large so that this is an isomorphism). Similarly,

$$\begin{aligned} H_k(\mathrm{Aut}(V); \mathbf{k}) &= H_k(\mathrm{Aut}(V \oplus X^m); \mathbf{k}[\mathrm{Hom}(X^m, V \oplus X^m)]) \\ &\cong H_k(\mathrm{Aut}(V \oplus X^m \oplus Y); \mathbf{k}[\mathrm{Hom}(X^m, V \oplus X^m \oplus Y)]) \\ &= H_k(\mathrm{Aut}(V \oplus Y); \mathbf{k}). \end{aligned}$$

These are enough to imply the conclusion. \square

Acknowledgements This paper is the result of PRIMES-USA, a program in which high school students (the second author) engage in research-level mathematics led by a mentor (the first author). This paper is based upon work supported by The National Science Foundation Graduate Research Fellowship Program under Grant No. 1842490 awarded to the first author. The authors would like to thank the organizers of the PRIMES-USA program for providing the opportunity for this research. We are also grateful to Dr. Tanya Khovanova and Dr. Kent Vashaw for their helpful comments and to Prof. Steven V. Sam for suggesting this research topic and providing valuable suggestions.

Funding Open Access funding provided by the MIT Libraries.

Declarations

Conflict of Interests Data sharing is not applicable to this article as no datasets were generated or analysed during the current study. The authors have no conflicts of interest to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Baeza, R.: Quadratic forms over semilocal rings, vol. 655. Springer (2006)
2. Brown, K.S.: Cohomology of Groups, vol. 87. Graduate texts in mathematics. Springer (1994)
3. Church, T.: Homological stability for configuration spaces of manifolds. In: Invent Math, vol. 188.2, pp. 465–504 (2012)
4. Church, T., Ellenberg, J.S., Farb, B.: FI-modules and stability for representations of symmetric groups. In: Duke Mathematical Journal, vol. 164.9, pp 1833–1910. issn: 0012-7094 (2015)
5. Church, T., Farb, B.: Representation theory and homological stability. In: Advances in Mathematics, vol. 245, pp. 250–314. issn: 0001-8708 (2013)
6. Friedlander, E.M.: Homological Stability for Classical Groups over Finite Fields. Lecture Notes in Mathematics, vol. 551. Springer (1976)
7. Gan, W.L., Li, L.: On central stability. arXiv:1504.07675 (2015)
8. Ganske, G., McDonald, B.R.: Finite local rings. In: The Rocky Mountain Journal of Mathematics, pp. 521–540 (1973)

9. Glasser, D.: Classification of bilinear forms over finite fields. <http://wwwmath.mit.edu/~dav/Fqquadforms.pdf>. Accessed 15 Feb 2021 (2005)
10. Lam, T.Y.: A first course in noncommutative rings. Graduate texts in mathematics. Springer, isbn: 9780387951836 (2001)
11. McDuff, D.: Configuration spaces of positive and negative particles. In: *Topology*, vol. 14, pp. 91–107 (1975)
12. Miller, J., Wilson, J.C.H.: Quantitative representation stability over linear groups. In: *International Mathematics Research Notices*, vol. 2020.22, pp. 8624–8672 (2020)
13. Putman, A., Sam, S.V.: Representation stability and finite linear groups. In: *Duke Mathematical Journal*, vol. 166.13, pp. 2521–2598. issn:0012-7094 (2017)
14. Sam, S.V., Snowden, A.: Gröbner methods for representations of combinatorial categories. In: *J. Amer. Math. Soc.* vol. 30, pp. 159–203 (2017)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.