



Association of short-term nitrogen dioxide exposure with hospitalization for urolithiasis in Xinxiang, China: a time series study

Yangdong Li¹ · Yongbin Wang² · Maochuan Fan¹ · Weisheng Li³ · Xiangzhen Meng¹ · Hao Zhou¹ · Shaohua Zhang¹ · Qifeng Dou¹

Received: 2 December 2022 / Accepted: 28 June 2023

© The Author(s) 2023

Abstract

Urolithiasis accounts for the highest incidence of all urologic-associated hospitalizations. However, few studies have explored the effect of nitrogen dioxide (NO₂) on hospitalizations for urolithiasis. We included 5956 patients with urolithiasis, collected daily meteorological and air pollution data between 2016 and 2021, and analyzed the associations between air pollutants and hospitalization, length of the hospital stay, and hospitalization costs attributable to urolithiasis. NO₂ exposure was associated with an increased risk of hospitalization for urinary tract stones. For each 10-μg/m³ increase and 1-day lag of NO₂, the maximum daily effect on the risk of hospitalization for urolithiasis was 1.020 (95% confidence interval [CI]: 1.001–1.039), and the cumulative effect peaked on lag day 4 (relative risk [RR]: 1.061; 95% CI: 1.003–1.122). Attribution scores and quantitative analysis revealed that the mean number of hospital days and mean hospital costs were 16 days and 21,164.39 RMB, respectively. Up to 5.75% of all urolithiasis hospitalizations were estimated to be attributable to NO₂, and the cost of NO₂-related urolithiasis hospitalizations reached approximately 3,430,000 RMB. Stratified analysis showed that NO₂ had a more sensitive impact on urolithiasis hospitalizations in women and in those aged ≥65 years. Notably, men and those younger than 65 years of age (exclude people aged 65) incurred more costs for urolithiasis hospitalizations. In the population level, the association between NO₂ and risk of urolithiasis hospitalization was more pronounced during the warm season. NO₂ can increase hospitalizations for urolithiasis for Xinxiang City residents, and there is a cumulative lag effect. Focusing on air pollution may have practical significance in terms of the prevention and control of urolithiasis.

Keywords Air pollution · Hospitalization · NO₂ · Time series analysis · Urolithiasis · Short-term exposure

Introduction

Urolithiasis is one of the most common and complex diseases of the urinary system worldwide, and its onset is related to age, sex, diet, and the environment. Currently several air pollutants, including particulate matter (PM_{2.5} and PM₁₀), nitrogen dioxide (NO₂), carbon monoxide (CO), ozone (O₃), and sulfur dioxide (SO₂), have been found to be associated with kidney disease. Nowadays, studies have confirmed a strong correlation between NO₂ concentration and mortality, which is consistent with the long-term impact of PM_{2.5} on mortality (Atkinson et al. 2018). With the rapid development of the economy, issues associated with the atmospheric environment are increasingly drawing the public's attention, particularly in recent years during which the frequent haze weather and environmental quality decline have caused a lot of adverse effects on people's

Responsible Editor: Lotfi Aleya

✉ Qifeng Dou
qifeng@263.net

¹ The First Affiliated Hospital of Xinxiang Medical University, No. 88, Jiankang Road, Weihui, Xinxiang, Henan Province 453100, People's Republic of China

² Department of Epidemiology and Health Statistics, School of Public Health, Xinxiang Medical University, Xinxiang, Henan Province 453003, People's Republic of China

³ The First Affiliated Hospital of Henan University of Chinese Medicine, Zhengzhou, Henan Province 450003, People's Republic of China

health. Moreover, industrial production processes, garbage incineration, and urban residential life are the main sources of NO₂, and people's lack of self-protection awareness from NO₂ pollution will also have a certain impact on human health (Zhu et al. 2019).

In addition, most of the major air pollutants are strong oxidants that induce oxidative stress in human cells, which is a modifiable risk factor for cell damage. NO₂ and O₃ are the two most important gaseous air pollutants, and both have strong oxidation capacity. Therefore, exposure to NO₂, O₃, and PM is associated with the occurrence of oxidative stress in human body (Zhang et al. 2021; Afsar et al. 2019; Mohammad et al. 2021). At present, the concentration of NO₂ in the atmosphere has become one of the indicators for measuring the strength of air pollution, which is associated with the occurrence and development of human diseases, such as chronic kidney (Wu et al. 2022), cardiovascular (Huang et al. 2021), and respiratory system (Liang et al. 2021) disease. However, little is known about the association between air pollutants and the risk of hospitalization for urinary stones. A better understanding of this association will be of great significance for the development of prevention and treatment strategies for urolithiasis.

Therefore, this study aimed to use the distributed hysteresis nonlinear model to analyze the nonlinear relationship and hysteresis effect between air pollutants and meteorological factors and the number of hospitalizations, number of hospitalization days, and hospitalization costs associated with urolithiasis in Xinxiang City, China, in order to provide a reference for reducing the influence of air pollutants on the risk of hospitalization of urolithiasis.

Materials and methods

Study area

This study was conducted in Xinxiang, China, which is located in the hinterland of the Central Plains in the northern part of Henan Province. The area has a warm temperate continental climate. Xinxiang is an important industrial base and central city in the Central Plains, and is a key city involved in the “2 + 26” air pollution monitoring program in the Beijing-Tianjin-Hebei region and surrounding areas.

Study population and data source

We collected data of a total of 5956 hospitalizations for urolithiasis from the First Affiliated Hospital of Xinxiang Medical College, the Third Affiliated Hospital of Xinxiang Medical College, Xinxiang Maternal and Child Health Hospital, and the First People's Hospital of Xinxiang City. The study period ranged from January 1, 2016 to October 31, 2021. The collected data

included the number of hospitalizations attributable to urolithiasis, average number of days in the hospital, date of hospitalization, date of discharge, age, sex, and medical costs during hospitalization. The 10th edition of the *International Classification of Diseases* (ICD-10) was used to statistically organize urolithiasis (ICD-10 code range: N20–N23). Meteorological data, including the average daily temperature, average daily relative humidity, and average daily barometric pressure during the study period, were obtained from the National Meteorological Science Center (<http://data.cma.cn/>). The average daily mass concentrations of air pollutants, such as PM_{2.5}, PM₁₀, NO₂, CO, SO₂, and the daily maximum 8-h mean concentrations of ambient O₃ during the study period, were obtained from the China Environmental Monitoring Station (<http://www.cnemc.cn>).

Theory/calculation

The SPSS software (IBM, SPSS Inc., New York, NY, USA) was used for data processing and analyses. Descriptive statistics, such as the mean, minimum, maximum, 25th percentile (P₂₅), 50th percentile (P₅₀), and 75th percentile (P₇₅), were used to describe the total number of hospitalizations for urolithiasis, the average length of the hospital stay (days), sex, age, meteorological factors, and air pollutants. The “DLNM,” “mgcv,” and “spline” packages of the R software (version 4.2.1) were used to construct the distributed lag nonlinear model (DLNM). The total number of hospitalizations for urolithiasis was considered as the dependent variable. A daily average temperature cross base was established, and the following were incorporated into the model: natural cubic spline function of long-term trends in pollutant data (PM_{2.5}, PM₁₀, NO₂, SO₂, CO, and O₃), average daily relative humidity, time, day of the week effect, and holiday effect. This resulted in the following equation (Wu et al. 2022):

$$\begin{aligned} \text{Logit}[E(Y_i)] = & \alpha + \beta \times X_{\text{pollutant}} + ns(\text{time}, df = 7 \times \text{year}) \\ & + ns(\text{temp}, df = 3) + ns(\text{humidity}, df = 3) \\ & + \text{dow} + \text{holidays} \end{aligned}$$

In the above formula, Y_i is the number of hospitalizations for urolithiasis, α is the intercept, β is the regression coefficient, $X_{\text{pollutant}}$ denotes the pollutant of interest, ns denotes the natural spline function, df is the degrees of freedom, $temp$ is the temperature, $humidity$ is the relative humidity, dow is the variable for the day of the week effect, and $holidays$ is the holiday effect.

We estimated the economic loss associated with NO₂ at the individual level for the attributable fraction (AF) and the attributable number (AN) of hospitalizations for urolithiasis, that is, the amount spent on urolithiasis per capita. Then, we evaluated the potential health and economic benefits of maintaining NO₂ concentrations below the current NO₂ air quality standard in China (24-h mean: 40 µg/m³) during the

study period (Guo et al. 2019) using the following equation (Miettinen 1974):

$$PAF = P(E|D) \times (RR - 1)/RR$$

In the above equation, $P(E|D)$ represents the proportion of exposure and disease in the population and RR represents the relative risk of exposure.

We stratified inpatients with urolithiasis according to season (warm season: May–October; cold season: November–April), age (younger than 64 years; 65 years or older), and sex. To assess the independence of the association of NO_2 exposure with urolithiasis, we fitted a dual-pollutant model by introducing other gaseous air pollutants, including $\text{PM}_{2.5}$, PM_{10} , SO_2 , CO , and O_3 . In this study, a DLNM was used to analyze the nonlinear relationship between, and the lag effects on the risk of, hospitalization for urolithiasis and air pollutants and meteorological factors.

We calculated the effect estimate and 95% confidence interval (CI) for the association between NO_2 exposure and urinary stones. $P < 0.05$ was considered statistically significant.

Results

Basic characteristics

A total of 5956 cases of urolithiasis were included in this study. Table 1 shows the descriptive statistics for urolithiasis, air pollutants, and meteorological factors during the study period. The median daily concentrations of $\text{PM}_{2.5}$, PM_{10} , NO_2 , CO , and SO_2 were $46.00 \mu\text{g}/\text{m}^3$, $94.00 \mu\text{g}/\text{m}^3$, $40.00 \mu\text{g}/\text{m}^3$, $1.00 \text{mg}/\text{m}^3$, and $16.00 \mu\text{g}/\text{m}^3$, respectively, and the maximum 8-h concentration of O_3 was $97.00 \mu\text{g}/\text{m}^3$. The average length of the hospital stay and average

Table 1 Descriptive statistics of hospitalizations for urolithiasis, length of the hospital stay, hospitalization costs, air pollutants, and meteorological factors in Xinxiang, China, from January 1, 2016 to October 31, 2021

	$\bar{x} \pm s$	Minimum	25th	50th	75th	Maximum
Hospitalizations (counts)						
Total	2.79±2.18	0.00	1.00	2.00	4.00	15.00
Male	1.85±1.64	0.00	1.00	2.00	3.00	10.00
Female	0.95±1.10	0.00	0.00	1.00	1.00	8.00
<65 years	2.48±1.991	0.00	1.00	2.00	4.00	14.00
≥65 years	0.310±0.593	0.00	0.00	0.00	1.00	4.00
Hospitalizations (stays in days)						
Total	20.38 ±18.93	0.00	6.00	16.00	30.00	181.00
Male	12.83±14.19	0.00	1.00	9.00	19.00	139.00
Female	7.55± 10.73	0.00	0.00	3.00	11.00	96.00
<65 years	17.62± 16.97	0.00	5.001	14.00	25.00	181.00
≥65 years	2.76 ±6.43	0.00	0.00	0.00	1.00	84.00
Hospitalizations costs (RMB)						
Total	28,012.87 ±29,394.88	0.00	3746.05	21,164.39	41,533.24	310,457.87
Male	17,747.00± 21,831.64	0.00	558.80	10,859.96	27,771.75	273,340.76
Female	10,265.87 ±16,318.32	0.00	0.00	1644.50	16,204.45	109,953.05
<65 years	24,151.21 ±26,469.59	0.00	2440.00	18,104.86	35,537.17	310,457.87
≥65 years	3861.66±9767.81	0.00	0.00	0.00	0.00	105,821.50
Meteorological factors						
Mean temperature (°C)	16.23± 10.11	−8.00	7.36	17.10	25.50	34.60
Relative humidity (%)	61.26 ±18.11	13.00	48.00	63.00	75.26	100.00
Air pollutants						
$\text{PM}_{2.5}$ ($\mu\text{g}/\text{m}^3$)	61.04±47.27	5.00	32.00	46.00	75.00	644.00
PM_{10} ($\mu\text{g}/\text{m}^3$)	111.70±71.46	12.00	66.00	94.00	136.50	823.00
NO_2 ($\mu\text{g}/\text{m}^3$)	42.32±19.94	8.00	27.00	40.00	55.00	168.00
CO (mg/m^3)	1.54± 0.70	0.30	0.70	1.00	1.37	8.00
SO_2 ($\mu\text{g}/\text{m}^3$)	21.31± 16.88	3.00	11.00	16.00	25.00	156.00
O_3 ($\mu\text{g}/\text{m}^3$)	104.20 ±55.01	7.00	60.00	97.00	145.00	276.00

hospitalization cost across all urolithiasis cases were 16 days and 21,164.39 RMB, respectively. The median daily temperature was 17.10 °C, and the median daily relative humidity was 63.00%.

A Spearman correlation analysis was performed to analyze the correlation between various variables of air pollutants and meteorological factors. These correlations are provided in Table 2.

In this study, from 2016 to October 31, 2021, there were 5956 hospitalizations for urolithiasis in Xinxiang, China. Of these hospitalizations, 3937 (66.10%) related to male and 2019 (33.90%) to female patients; therefore, the male-to-female ratio was approximately 1.95 (see Table 1). A time series plot of the number of hospitalizations for urolithiasis, length of the hospital stay, and hospitalization costs (Fig. 1; Figure S1) showed that the median number of hospitalizations per day was two. The total number of hospitalizations for urolithiasis was 5287 (88.77%) for those 65 years of age or younger. However, the total number of hospitalizations for urolithiasis was 669 (11.23%) for those older than 65 years of age.

Effects of six air pollutants on hospitalizations for urinary tract stones

Using the single-pollutant model (see Table 3), we observed that the number of hospitalizations for urolithiasis increased with higher NO₂ concentrations. Furthermore, it was observed that NO₂ had a nonlinear relationship with the number of hospitalizations for urolithiasis (Fig. 2; Figure S2). The maximum single-day effect of NO₂ on hospitalizations for urolithiasis was 1.020 (95% CI: 1.001–1.039 with a 1-day lag) for each 10-μg/m³ increase in the NO₂ concentration. The cumulative effect on hospitalizations for urolithiasis peaked on lag day 4, with a relative risk (RR) of 1.061 (95% CI: 1.003–1.122). The results for the other pollutants are summarized in Fig. 2 and Figure S2.

Table 4 summarizes the results of the two-pollutant model for air pollutants. After adjusting for PM_{2.5}, PM₁₀, SO₂, CO, and O₃, the association between NO₂ exposure (lag day 1) and hospitalizations for urolithiasis remained robust. Moreover, the correlation between NO₂ (cumulative lag days 2, 3, and 4) and hospitalizations for urolithiasis was significantly enhanced after controlling for PM_{2.5}, PM₁₀, SO₂, CO, and O₃.

Table 2 Spearman correlation between daily meteorological factors and air pollutants

Pollutants	PM _{2.5}	PM ₁₀	CO	NO ₂	SO ₂	O ₃	Temp	RH
PM _{2.5}	1.000							
PM ₁₀	0.869**	1.000						
CO	0.750**	0.612**	1.000					
NO ₂	0.683**	0.723**	0.620**	1.000				
SO ₂	0.476**	0.583**	0.497**	0.622**	1.000			
O ₃	-0.400**	-0.329**	-0.415**	-0.409**	-0.114**	1.000		
Temp	-0.532**	-0.493**	-0.456**	-0.488**	-0.264**	0.800**	1.000	
RH	0.126**	-0.172**	0.178**	-0.154**	-0.457**	-0.145**	0.119**	1.000

Temp, temperature; RH, relative humidity

**P-value < 0.001

Fig. 1 Time series of hospitalizations for urolithiasis in Xinxiang, China, from January 1, 2016 to October 31, 2021




Table 3 Single-day and cumulative lag effects of 10- $\mu\text{g}/\text{m}^3$ increases in the concentrations of six pollutants on the number of hospitalizations for urolithiasis in Xinxiang, China, from January 1, 2016 to October 31, 2021

	PM _{2.5}		PM ₁₀		NO ₂		SO ₂		CO		O ₃	
	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI
Lag 0	1.012	(0.980–1.053)	0.999	(0.988–1.011)	1.014	(0.976–1.053)	0.993	(0.948–1.040)	1.025	(0.958–1.097)	0.991	(0.976–1.007)
Lag 1	1.016	(1.001–1.039)	1.000	(0.995–1.006)	1.020	(1.001–1.039)	1.006	(0.984–1.029)	1.002	(0.972–1.034)	1.005	(0.997–1.013)
Lag 2	1.014	(0.996–1.039)	1.000	(0.994–1.007)	1.017	(0.995–1.039)	1.010	(0.985–1.035)	0.993	(0.959–1.029)	1.009	(1.001–1.017)
Lag 3	1.008	(0.994–1.026)	1.000	(0.995–1.005)	1.009	(0.992–1.026)	1.008	(0.989–1.028)	0.994	(0.967–1.022)	1.008	(1.001–1.014)
Lag 4	1.000	(0.986–1.017)	1.000	(0.995–1.005)	1.000	(0.983–1.017)	1.003	(0.984–1.023)	1.000	(0.973–1.028)	1.003	(0.997–1.009)
Lag 5	0.994	(0.976–1.014)	1.001	(0.994–1.007)	0.993	(0.972–1.014)	0.999	(0.975–1.024)	1.007	(0.972–1.044)	0.999	(0.991–1.006)
Lag 6	0.992	(0.977–1.009)	1.002	(0.996–1.008)	0.991	(0.973–1.009)	0.998	(0.977–1.020)	1.012	(0.982–1.042)	0.998	(0.991–1.005)
Lag 7	0.998	(0.968–1.034)	1.005	(0.994–1.016)	0.997	(0.962–1.034)	1.004	(0.964–1.046)	1.009	(0.950–1.071)	1.004	(0.991–1.017)
Lag 0–1	1.028	(0.990–1.068)	0.999	(0.986–1.014)	1.034	(0.988–1.082)	0.999	(0.944–1.057)	1.027	(0.948–1.114)	0.996	(0.976–1.016)
Lag 0–2	1.043	(1.001–1.086)	1.000	(0.985–1.015)	1.051	(1.001–1.105)	1.009	(0.949–1.073)	1.021	(0.937–1.111)	1.005	(0.983–1.028)
Lag 0–3	1.050	(1.005–1.098)	1.000	(0.984–1.017)	1.061	(1.006–1.119)	1.017	(0.952–1.087)	1.015	(0.926–1.111)	1.013	(0.989–1.037)
Lag 0–4	1.050	(1.002–1.100)	1.000	(0.983–1.018)	1.061	(1.003–1.122)	1.021	(0.952–1.096)	1.015	(0.921–1.118)	1.016	(0.992–1.041)
Lag 0–5	1.044	(0.992–1.098)	1.001	(0.981–1.021)	1.053	(0.990–1.119)	1.020	(0.944–1.102)	1.022	(0.917–1.139)	1.015	(0.989–1.041)
Lag 0–6	1.036	(0.979–1.095)	1.003	(0.981–1.026)	1.043	(0.975–1.115)	1.018	(0.935–1.109)	1.034	(0.916–1.167)	1.013	(0.986–1.040)
Lag 0–7	1.033	(0.975–1.095)	1.008	(0.984–1.032)	1.040	(0.970–1.115)	1.022	(0.935–1.118)	1.043	(0.919–1.183)	1.016	(0.990–1.044)

Fig. 2 Cumulative effect of NO₂ exposure on the number of hospitalizations for urolithiasis on different lag days




Table 4 Single-day and cumulative lag effects of 10- $\mu\text{g}/\text{m}^3$ increases in NO₂ and other air pollutant concentrations (PM_{2.5}, PM₁₀, SO₂, CO, and O₃) on hospitalizations for urolithiasis in a two-pollutant model from January 1, 2016 to October 31, 2021

	NO ₂ + PM _{2.5}		NO ₂ + PM ₁₀		NO ₂ + SO ₂		NO ₂ + CO		NO ₂ + O ₃	
	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI
Lag 0	1.021	(0.974–1.071)	1.022	(0.977–1.069)	1.028	(0.980–1.078)	1.011	(0.963–1.063)	1.019	(0.981–1.059)
Lag 1	1.041	(1.016–1.067)	1.033	(1.009–1.057)	1.027	(1.003–1.051)	1.035	(1.010–1.061)	1.020	(1.002–1.040)
Lag 2	1.040	(1.012–1.068)	1.029	(1.002–1.057)	1.019	(0.992–1.046)	1.036	(1.008–1.066)	1.015	(0.994–1.037)
Lag 3	1.024	(1.002–1.047)	1.016	(0.995–1.038)	1.007	(0.986–1.029)	1.022	(1.000–1.045)	1.007	(0.990–1.024)
Lag 4	1.002	(0.980–1.024)	1.000	(0.979–1.021)	0.996	(0.975–1.018)	1.002	(0.980–1.025)	0.999	(0.982–1.016)
Lag 5	0.981	(0.954–1.007)	0.985	(0.960–1.011)	0.987	(0.961–1.014)	0.985	(0.957–1.013)	0.992	(0.971–1.014)
Lag 6	0.967	(0.944–0.989)	0.977	(0.955–0.999)	0.985	(0.962–1.007)	0.977	(0.954–1.001)	0.991	(0.973–1.009)
Lag 7	0.966	(0.923–1.011)	0.979	(0.938–1.023)	0.991	(0.947–1.037)	0.988	(0.943–1.035)	0.997	(0.961–1.034)
Lag 0–1	1.063	(1.002–1.128)	1.056	(0.999–1.115)	1.056	(0.997–1.118)	1.047	(0.986–1.112)	1.040	(0.994–1.089)
Lag 0–2	1.105	(1.036–1.179)	1.086	(1.022–1.154)	1.076	(1.012–1.143)	1.085	(1.018–1.157)	1.056	(1.005–1.110)
Lag 0–3	1.132	(1.055–1.215)	1.104	(1.033–1.180)	1.084	(1.015–1.157)	1.109	(1.035–1.188)	1.064	(1.009–1.123)
Lag 0–4	1.134	(1.053–1.221)	1.103	(1.029–1.184)	1.079	(1.008–1.155)	1.111	(1.035–1.193)	1.063	(1.004–1.124)
Lag 0–5	1.112	(1.027–1.204)	1.087	(1.008–1.172)	1.065	(0.990–1.147)	1.094	(1.014–1.181)	1.054	(0.991–1.121)
Lag 0–6	1.075	(0.987–1.170)	1.062	(0.978–1.152)	1.049	(0.969–1.136)	1.070	(0.986–1.160)	1.044	(0.976–1.117)
Lag 0–7	1.039	(0.952–1.133)	1.040	(0.955–1.133)	1.039	(0.958–1.128)	1.057	(0.975–1.146)	1.041	(0.970–1.117)

Attributable fraction and number

Other measures such as disease burden (attributable score and number), health burden (length of stay), and economic burden (cost of stay) were used to further understand the effect of NO₂ on urolithiasis. In this study, we estimated that up to 5.75% of the total hospitalizations for urolithiasis could be attributed to NO₂, and the cost of hospitalizations for urolithiasis related to NO₂ amounted to approximately RMB 343 million (Table 5). It is worth noting that the magnitude of hospitalization risk for NO₂ varies by sex, age, and season, indicating that the resulting burden of disease is significantly comparable. A bar chart was also used to analyze the distribution of continuous variables for hospitalizations, hospital stays, and hospitalization costs (supplementary Figure S3). Other measures such as disease burden (attributable score and number), health burden (length of stay), and economic burden (cost of stay) were used to further understand the effect of NO₂ on urolithiasis disease.

Effect of increased NO₂ exposure on the number of hospitalizations for urolithiasis after stratification based on age, sex, and season

The results of the age-stratified analysis (see Fig. 3) showed that for every 10- $\mu\text{g}/\text{m}^3$ increase in the NO₂ concentration, the correlation between age 65 years or older and the number of hospitalizations for urolithiasis was high. Additionally, the maximum single-day effect was 1.074 (95% CI: 1.012–1.139) on lag day 2, and the cumulative effect peaked at 1.234 (95% CI: 1.054–1.445) on lag day 4 (detailed results of the hierarchical analysis are shown in Table S1).

The results of the sex-stratified analysis (see Fig. 4) showed that for every 10- $\mu\text{g}/\text{m}^3$ increase in the NO₂ concentration, the correlation between female patients and hospitalizations for urolithiasis was high. Additionally, the maximum single-day effect was 1.033 (95% CI: 1.001–1.065) on lag day 1, and the cumulative effect peaked at 1.100 (95% CI: 1.008–1.201)

on lag day 3 (detailed results of the hierarchical analysis are provided in Table S2).

The results of the seasonal stratification analysis (see Fig. 5) showed that the correlation between the warm season and the number of hospitalizations for urolithiasis was greater for each 10- $\mu\text{g}/\text{m}^3$ increase in the NO_2 concentration when the season was divided into cold and warm seasons. Additionally, the maximum single-day effect was 1.065 (95% CI: 1.020–1.113)

on lag day 2, and the cumulative effect peaked at 1.202 (95% CI: 1.039–1.391) on lag day 4 (detailed results of the stratified analysis are provided in Table S3).

Model stability analysis

By adjusting the degree of freedom of the natural spline function of time (8–13 df), it can be found that the relative

Table 5 Fraction and number of urolithiasis hospitalizations, hospital stays, and hospitalization costs attributable to NO_2 from January 1, 2016 to October 31, 2021

	Attributable fraction of hospitalizations (%)	Attributable number of hospitalizations (no.)	Attributable number of hospital stays (days)	Attributable number of hospitalization costs (RMB, million)
Total	5.75 (0.60–10.63)	342.47 (35.74–633.12)	2497.40 (260.60–4616.93)	3.43 (0.36–6.35)
Males	4.40 (–2.35–10.63)	173.23 (–92.52–418.50)	1203.14 (–642.58–2906.67)	1.66 (–0.89–4.02)
Females	3.19 (0.10–6.10)	64.41 (2.02–123.16)	513.24 (16.09–981.43)	0.70 (0.02–1.33)
Young (<65)	4.58 (–1.01–9.83)	242.14 (–53.40–519.71)	1719.38 (–379.16–3690.28)	2.36 (–0.52–5.06)
Elderly (≥ 65)	18.96 (5.12–30.80)	126.84 (34.25–206.05)	1117.12 (301.67–1814.74)	1.56 (0.42–2.53)
Warm	16.81 (3.75–28.11)	487.99 (108.86–816.03)	3550.27 (792.00–5936.83)	4.96 (1.11–8.30)
Cold	1.19 (–6.72–8.51)	36.33 (–205.16–259.81)	265.52 (–1499.43–1898.84)	0.36 (–2.03–2.57)

Note that figures in the parenthesis were the 95% confidence interval

Fig. 3 Hysteresis and cumulative hysteresis effects of each 10- $\mu\text{g}/\text{m}^3$ increase in the NO_2 concentration on the number of hospitalizations for urolithiasis on different lag days in the age-stratified analysis




Fig. 4 Hysteresis and cumulative hysteresis effects of each 10- $\mu\text{g}/\text{m}^3$ increase in the NO_2 concentration on the number of hospitalizations for urolithiasis on different lag days in the sex-stratified analysis





Fig. 5 Lag and cumulative lag effects of each $10\text{-}\mu\text{g}/\text{m}^3$ increase in the NO_2 concentration on the number of hospitalizations for urolithiasis on different lag days in the seasonal stratification analysis



risk of the model has no obvious change, so it can be considered that the established model has stability, and the results of model analysis are reliable (Table 6).

Discussion

This study used the DLNM model to investigate the association between air pollutants and daily mean temperature on the number of hospitalizations for urolithiasis, hospital length of stay, and hospitalization costs in Xinxiang City, China, from January 1, 2016 to October 31, 2021. The results showed a significant correlation between short-term NO_2 exposure and the number of hospitalizations for urolithiasis. The correlation was more significant for elderly patients, female patients, and warm seasons. The association between NO_2 exposure and urolithiasis was robust even after adjusting for other air pollutants.

Based on the DLNM analysis results, our study demonstrated that the number of daily hospitalizations in the time series presented a periodic distribution with the cold and warm seasons, which was higher in the warm (May to September) and lower in the cold (October to April of the next year) season. The concentration of various air pollutants also presented a cyclical trend, with O_3 , $\text{PM}_{2.5}$, PM_{10} , NO_2 , SO_2 , and CO being cyclically opposite with temperature, mainly high in the warm season and low in the cold season. In the single-pollutant model analysis, there was a nonlinear relationship between NO_2 and the number of inpatients with urolithiasis, and there was a certain lag effect. This study found that NO_2 exposure was associated with an increased risk of hospitalization for urolithiasis. At daily mean temperature, the maximum daily effect of NO_2 and $\text{PM}_{2.5}$ increased by $10\text{ }\mu\text{g}/\text{m}^3$ on the hospitalization risk of urolithiasis was 1 day behind, and the cumulative lag effect reached a peak on day 4. When O_3 increased by $10\text{ }\mu\text{g}/\text{m}^3$, the maximum 1-day effect of O_3 on the number of hospitalizations for urolithiasis was 2 days behind. However, no cumulative hysteretic effect of O_3 on the number of hospitalizations for urolithiasis was found. No other association was observed between the concentrations

of other pollutants (PM_{10} , SO_2 , and CO) and the number of hospitalizations for urolithiasis per increase of $10\text{ }\mu\text{g}/\text{m}^3$ or $10\text{ mg}/\text{m}^3$. These results suggest that short-term NO_2 exposure is a potential cause of increased risk of hospitalization for urolithiasis. In recent years, air pollution is an important factor causing kidney disease, mainly because the kidney is susceptible to the toxic effects of air pollutants (Xu et al. 2018). One study (Chen et al. 2018) found that NO_2 was associated with a lower glomerular filtration rate and a higher prevalence of chronic kidney disease. Existing studies have also confirmed that NO_2 is a major atmospheric pollutant affecting the hospitalization rate of chronic kidney disease (Wu et al. 2020; Ye et al. 2021). In addition, NO_2 is correlated to the occurrence of oxidative stress. More and more studies have confirmed that oxidative stress is strongly correlated with the formation of kidney stones (Jeong et al. 2020; Wang et al. 2022), mainly because it can maintain an inflammatory microenvironment, reshape the extracellular matrix, and lead to the accumulation of crystal deposits, thus promoting kidney stone formation (Wigner et al. 2021). In conclusion, although the exact biological pathways by which air pollution induces kidney injury are not fully understood, it is possible that air pollution-related renal toxicity is caused by the same biological mechanisms through which air pollution exerts a negative impact on the cardiovascular system, including oxidative stress and systemic inflammation, blood pressure changes, or vascular damage (Rasking et al. 2022). Therefore, the correlation between NO_2 and urolithiasis needs to be further explored.

Our dual-pollutant model demonstrated that after adjusting for $\text{PM}_{2.5}$, PM_{10} , SO_2 , CO , and O_3 , the association between NO_2 (day 1 lag) and number of hospitalizations for urolithiasis remained robust. In addition, $\text{PM}_{2.5}$, PM_{10} , SO_2 , CO , and O_3 could increase the risk of hospitalization of urolithiasis when the cumulative lag was 3 days. This further suggests a strong association between NO_2 exposure and hospitalizations for urolithiasis. It has been reported that ambient PM and NO_2 may contribute to kidney damage through peroxidation. It has also been shown that factors affecting the formation of urolithiasis include $\text{PM}_{2.5}$, potentially by affecting serum creatinine and uric acid levels (Aizezi et al. 2022). In addition, studies have shown

Table 6 Sensitivity analysis result

df	Temperature degree of freedom	Male		Female		Young (<65)		Elderly (≥65)		Warm		Cold	
		RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI
		3	1.013	(0.990–1.037)	1.033	(1.001–1.066)	1.013	(0.993–1.034)	1.075	(1.019–1.134)	1.050	(1.006–1.095)	1.004
3	1.011	(0.988–1.035)	1.028	(0.996–1.061)	1.011	(0.990–1.032)	1.072	(1.016–1.132)	1.046	(1.002–1.092)	0.999	(0.968–1.031)	
3	1.014	(0.990–1.038)	1.028	(0.995–1.062)	1.011	(0.990–1.032)	1.084	(1.025–1.146)	1.049	(1.003–1.096)	0.993	(0.961–1.026)	
3	1.008	(0.985–1.032)	1.024	(0.991–1.057)	1.006	(0.986–1.027)	1.074	(1.017–1.137)	1.050	(1.004–1.097)	0.997	(0.962–1.035)	
3	1.009	(0.986–1.034)	1.025	(0.993–1.059)	1.008	(0.987–1.029)	1.077	(1.019–1.139)	1.047	(1.000–1.095)	0.993	(0.954–1.032)	
3	1.009	(0.986–1.034)	1.023	(0.990–1.057)	1.008	(0.987–1.029)	1.072	(1.013–1.134)	1.056	(1.006–1.107)	1.008	(0.966–1.052)	

that O₃ exposure is related to elevated levels of systemic inflammation (Arjomandi et al. 2015), related to the up-regulation of redox homeostasis in vivo caused by long-term continuous exposure to O₃ (Hu et al. 2020). Moreover, it has been shown that high concentration of environmental PM and O₃ exposure could lead to inflammation and oxidative stress in the perirenal adipose tissue of rats (Sun et al. 2013). Finally, an experimental study in mice exposed to PM_{2.5} for a long time revealed increased levels of DNA damage related to renal oxidative stress (Bowe et al. 2020). Therefore, based on previous studies, we speculate that the effect of air pollutants on urolithiasis may be related to increased levels of oxidative stress and inflammation in the body. Future studies should further explore this pathogenic mechanism.

According to the attribution score and quantitative analysis, the average length and the average cost of hospitalization were 16 days and 21,164.39 RMB, respectively. It is estimated that up to 5.75% of the total hospitalizations for urolithiasis can be attributed to NO₂, and the cost of hospitalizations for urolithiasis related to NO₂ is about 3,430,000 RMB. In addition, there is a large difference in the incidence rate of urolithiasis between male and female, with women having a significantly higher incidence rate compared to men. However, it is worth noting that even though the prevalence rate of female is higher than that of male, the number of hospitalizations, length of stay, and hospitalization cost of men are higher than those of women. This result indicates that men have a certain economic burden and human and material loss due to urolithiasis. Similarly, in terms of age groups, we concluded that the prevalence of urolithiasis was higher in those aged ≥65 years compared to people aged <65 years, but the number of hospital days, hospital admissions, and hospital costs after urolithiasis were greater in those aged <65 years compared to those aged ≥65 years. These findings suggest that people younger than 65 spend more time, energy, and money on urolithiasis. In addition, different populations are more susceptible to urolithiasis during warm seasons than during cold seasons. Therefore, based on in this study, the relevant government departments should strengthen health education on urolithiasis to improve the public’s awareness of the disease. At the same time, the relevant environmental departments should also strengthen public awareness regarding environmental protection, in order to reduce the morbidity and economic losses caused by environmental pollution.

We conducted a stratified analysis based on age, sex, and season. The results showed that with increasing NO₂ concentrations, age 65 years or older was more strongly correlated with hospitalizations for urolithiasis than age younger than 65 years. In terms of age effects, elderly individuals (older than 65 years of age) are more susceptible to air pollutants than other age groups because of their weaker immune systems (Tong et al. 2016). The results of this study showed that the risk of hospitalization for urolithiasis increases with increasing NO₂ concentrations, and this effect was stronger for females than for males. This finding is consistent with that of another study

(Safdar et al. 2021). Other studies have also shown that females and individuals older than 65 years of age seem to be more susceptible to the effects of air pollution than males and younger individuals (Manisalidis et al. 2020). Previous studies have suggested that males are 2 to 3 times more likely to develop stones than females; however, data from recent studies suggest that this difference is decreasing (Khan et al. 2016). The adverse effects of air pollution and their association with sex differences have rarely been the focus of scholarly research. Sex differences in terms of behavior, occupation, and social/family roles cause differences in air pollution exposure, which may explain the variability in the risk associated with air pollution exposure of males and females (Matz et al. 2014). However, the current limited understanding of biological mechanisms presents a challenge when explaining the association between sex differences and air pollution in terms of its effect on the risk of hospitalization for urolithiasis. Additionally, our study found a stronger correlation between warm seasons and hospitalizations for urolithiasis with increasing NO₂ concentrations compared to cold seasons. Human sensitivity to air pollutants may vary with seasons because of physiological differences in hormones, biology, and structure/morphology. Another study (Shin et al. 2022) found seasonal differences in hospitalizations for circulatory and respiratory diseases and mortality rates in response to short-term exposure to air pollutants. However, reports related to urolithiasis are scarce. Therefore, more data from different regions are necessary to further explore the correlation between age, sex, and season, and the number of hospitalizations for urolithiasis as air pollutant concentrations increase.

The study was stratified by age, sex, and season, which could help target specific interventions to sensitive groups. However, several limitations of our study should be noted. First, air pollution data were obtained from fixed site monitors. Although errors are inevitable, the resulting nondifferential errors may have resulted in the underestimation of the impact of air pollutants (Richmond and Long 2020). Second, this study focused on the average daily temperature. Further studies should consider different temperature indicators, such as extreme and daily maximum temperatures. Finally, the relatively low daily number of urolithiasis cases may have caused us to miss important associations between air pollutants and urolithiasis, especially in subgroup analyses. Future multicenter studies with longer durations are required to validate these results.

Conclusions

Taken together, the results of this study suggest that short-term exposure to NO₂ is associated with an increased risk of hospitalization for urolithiasis. Among these hospitalizations, women and the elderly are more susceptible to air pollution, especially NO₂. Furthermore, the number of hospitalizations for

urolithiasis is higher during the warm season. Finally, men and people younger than 65 years of age incur more costs for urolithiasis-related hospitalizations. Further studies are needed to confirm our findings and the underlying mechanisms involved.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11356-023-28539-0>.

Acknowledgements The authors acknowledge the dedication and cooperation of all the participants in this study.

Author contribution Yangdong Li: conceptualization, methodology, formal analysis, investigation, resources, data curation, writing—original draft, writing—review and editing, and visualization. Weisheng Li: data curation, investigation and visualization, and formal analysis. Maochuan Fan: data curation, writing—review and editing, and visualization. Xiangzhen Meng, Hao Zhou, and Shaohua Zhang: data curation and data analysis. Yongbin Wang and Qifeng Dou: conceptualization, methodology, formal analysis, writing—review and editing, and read and approved the final version.

Funding This work was supported by Xinxiang Science and Technology Tackling Project (grant no. GG2020032), the Key Scientific Research Project of Henan Higher Education Institution (grant no. 21A320015), Xinxiang Post-Disaster Reconstruction Science and Technology Special Project (grant no. 21CJ002), and 2020 Youth Cultivation Fund of the First Affiliated Hospital of Xinxiang Medical University (grant no. QN-2020-A05).

Data availability The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethical approval Not applicable.

Consent to participate Not applicable.

Consent for publication All authors declare that they have consent for publication in the journal of Environmental Science and Pollution Research.

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Afsar B, Elsurur Afsar R, Kanbay A, Covic A, Ortiz A, Kanbay M (2019) Air pollution and kidney disease: review of current evidence. *Clin Kidney J* 12(1):19–32. <https://doi.org/10.1093/ckj/sfy111>

- Atkinson RW, Butland BK, Anderson HR, Maynard RL (2018) Long-term concentrations of nitrogen dioxide and mortality: a meta-analysis of cohort studies. *Epidemiology* 29:460–472. <https://doi.org/10.1097/EDE.0000000000000847>
- Aizezi X, Xie L, Xie H, Li J, Shang Z, Liu C (2022) Epidemiological and clinical characteristics of stone composition: a single-center retrospective study. *Urolithiasis* 50:37–46. <https://doi.org/10.1007/s00240-021-01274-2>
- Arjomandi M, Wong H, Donde A, Frelinger J, Dalton S, Ching W, Power K, Balmes JR (2015) Exposure to medium and high ambient levels of ozone causes adverse systemic inflammatory and cardiac autonomic effects. *American journal of physiology. Am J Physiol Heart Circ Physiol* 308:H1499–H1509. <https://doi.org/10.1152/ajpheart.00849.2014>
- Bowe B, Artimovich E, Xie Y, Yan Y, Cai M, Al-Aly Z (2020) The global and national burden of chronic kidney disease attributable to ambient fine particulate matter air pollution: a modelling study. *BMJ Global Health* 5(3):e002063. <https://doi.org/10.1136/bmjgh-2019-002063>
- Chen SY, Chu DC, Lee JH, Yang YR, Chan CC (2018) Traffic-related air pollution associated with chronic kidney disease among elderly residents in Taipei City. *Environ Pollut* 234:838–845. <https://doi.org/10.1016/j.envpol.2017.11.084>
- Guo H, Gu X, Ma G, Shi S, Wang W, Zuo X, Zhang X (2019) Spatial and temporal variations of air quality and six air pollutants in China during 2015–2017. *Sci Rep* 9:15201. <https://doi.org/10.1038/s41598-019-50655-6>
- Huang Y, Zhu M, Ji M, Fan J, Xie J, Wei X, Jiang X, Xu J, Chen L, Yin R, Wang Y, Dai J, Jin G, Xu L, Hu Z, Ma H, Shen H (2021) Air pollution, genetic factors, and the risk of lung cancer: a prospective study in the UK Biobank. *Am J Respir Crit Care Med* 204:817–825. <https://doi.org/10.1164/rccm.202011-4063OC>
- Hu X, He L, Zhang J, Qiu X, Zhang Y, Mo J, Day DB, Xiang J, Gong J (2020) Inflammatory and oxidative stress responses of healthy adults to changes in personal air pollutant exposure. *Environ Poll (Barking, Essex: 1987)* 263(Pt A):114503. <https://doi.org/10.1016/j.envpol.2020.114503>
- Jeong SM, Park JH, Kim HJ, Kwon H, Hwang SE (2020) Effects of abdominal obesity on the association between air pollution and kidney function. *Int J Obes* 44(7):1568–1576. <https://doi.org/10.1038/s41366-020-0540-8>
- Manisalidis I, Stavropoulou E, Stavropoulos A, Bezirtzoglou E (2020) Environmental and health impacts of air pollution: a review. *Front Public Health* 8:14. <https://doi.org/10.3389/fpubh.2020.00014>
- Khan SR, Pearle MS, Robertson WG, Gambaro G, Canales BK, Doizi S, Traxer O, Tiselius HG (2016) Kidney stones. *Nat Rev Dis Primers* 2:16008. <https://doi.org/10.1038/nrdp.2016.8>
- Liang S, Sun C, Liu C, Jiang L, Xie Y, Yan S, Jiang Z, Qi Q, Zhang A (2021) The influence of air pollutants and meteorological conditions on the hospitalization for respiratory diseases in Shenzhen City, China. *Int J Environ Res Public Health* 18(10):5120. <https://doi.org/10.3390/ijerph18105120>
- Mohammad KN, Chan EYY, Lau SY, Lam HCY, Goggins WB, Chong KC (2021) Relationship between acute kidney injury, seasonal influenza, and environmental factors: a 14-year retrospective analysis. *Environ Int* 153:106521. <https://doi.org/10.1016/j.envint.2021.106521>
- Miettinen OS (1974) Proportion of disease caused or prevented by a given exposure, trait or intervention. *Am J Epidemiol* 99:325–332. <https://doi.org/10.1093/oxfordjournals.aje.a121617>
- Matz CJ, Stieb DM, Davis K, Egyed M, Rose A, Chou B, Brion O (2014) Effects of age, season, gender and urban-rural status on time-activity: Canadian Human Activity Pattern Survey 2 (CHAPS 2). *Int J Environ Res Public Health* 11:2108–2124. <https://doi.org/10.3390/ijerph110202108>
- Rasking L, Vanbrabant K, Bové H, Plusquin M, De Vusser K, Roels HA, Nawrot TS (2022) Adverse effects of fine particulate matter on human kidney functioning: a systematic review. *Environ Health* 21(1):24. <https://doi.org/10.1186/s12940-021-00827-7>
- Richmond-Bryant J, Long TC (2020) Influence of exposure measurement errors on results from epidemiologic studies of different designs. *J Expo Sci Environ Epidemiol* 30(3):420–429. <https://doi.org/10.1038/s41370-019-0164-z>
- Safdar OY, Alblowi SS, Aboulola NA, Alharazy DT (2021) Renal stones and risk factors in Jeddah and Riyadh. *Saudi J Kidney Dis Transpl* 32:191–198. <https://doi.org/10.4103/1319-2442.318523>
- Sun L, Liu C, Xu X, Ying Z, Maiseyeu A, Wang A, Allen K, Lewandowski RP, Bramble LA, Morishita M, Wagner JG, Dvonch J, Sun Z, Yan X, Brook RD, Rajagopalan S, Harkema JR, Sun Q, Fan Z (2013) Ambient fine particulate matter and ozone exposures induce inflammation in epicardial and perirenal adipose tissues in rats fed a high fructose diet. *Part Fibre Toxicol* 10:43. <https://doi.org/10.1186/1743-8977-10-43>
- Shin HH, Maquiling A, Thomson EM, Park IW, Stieb DM, Dehghani P (2022) Sex-difference in air pollution-related acute circulatory and respiratory mortality and hospitalization. *Sci Total Environ* 806:150515. <https://doi.org/10.1016/j.scitotenv.2021.150515>
- Tong L, Li K, Zhou Q (2016) Season, sex and age as modifiers in the association of psychosis morbidity with air pollutants: a rising problem in a Chinese metropolis. *Sci Total Environ* 541:928–933. <https://doi.org/10.1016/j.scitotenv.2015.09.066>
- Wang C, Tsai JD, Wan L, Lin CL, Wei CC (2022) Association between gaseous air pollutants and idiopathic nephrotic syndrome in children: a 12-year population-based cohort study. *Ital J Pediatr* 48(1):70. <https://doi.org/10.1186/s13052-022-01269-8>
- Wu MY, Lo WC, Chao CT, Wu MS, Chiang CK (2020) Association between air pollutants and development of chronic kidney disease: a systematic review and meta-analysis. *Sci Total Environ* 706:135522. <https://doi.org/10.1016/j.scitotenv.2019.135522>
- Wigner P, Grębowski R, Bijak M, Szemraj J, Saluk-Bijak J (2021) The molecular aspect of nephrolithiasis development. *Cells* 10:1926. <https://doi.org/10.3390/cells10081926>
- Wu J, Ye Q, Fang L, Deng L, Liao T, Liu B, Lv X, Zhang J, Tao J, Ye D (2022) Short-term association of NO₂ with hospital visits for chronic kidney disease and effect modification by temperature in Hefei, China: a time series study. *Ecotoxicol Environ Saf* 237:113505. <https://doi.org/10.1016/j.ecoenv.2022.113505>
- Xu X, Nie S, Ding H, Hou FF (2018) Environmental pollution and kidney diseases. *Nat Rev Nephrol* 14:313–324. <https://doi.org/10.1038/nrneph.2018.11>
- Ye JJ, Wang SS, Fang Y, Zhang XJ, Hu CY (2021) Ambient air pollution exposure and risk of chronic kidney disease: a systematic review of the literature and meta-analysis. *Environ Res* 195:110867. <https://doi.org/10.1016/j.envres.2021.110867>
- Zhang Y, Liu D, Liu Z (2021) Fine particulate matter (PM_{2.5}) and chronic kidney disease. *Rev Environ Contam Toxicol* 254:183–215. https://doi.org/10.1007/398_2020_62
- Zhu Y, Zhan Y, Wang B, Li Z, Qin Y, Zhang K (2019) Spatiotemporally mapping of the relationship between NO₂ pollution and urbanization for a megacity in Southwest China during 2005–2016. *Chemosphere* 220:155–162. <https://doi.org/10.1016/j.chemosphere.2018.12.095>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.